

Heidi Lilja

Searching for Genes Predisposing to Common Dyslipidemias

Department of Molecular Medicine
National Public Health Institute, Helsinki, Finland
and
Medical Faculty,
University of Helsinki, Finland

2004



SEARCHING FOR GENES PREDISPOSING TO COMMON DYSLIPIDEMIAS

Heidi Lilja

Department of Molecular Medicine, National Public Health Institute, and Department of
Medical Genetics, University of Helsinki, and Department of Medicine,
Division of Cardiology, University of Helsinki,
Helsinki, Finland

Academic Dissertation

To be publicly discussed by the permission of the Medical Faculty of the University of
Helsinki, in the Lecture Hall 3 of Biomedicum, Helsinki,
on December 17th, at 12 noon.

Helsinki 2004

Supervised by:

Assistant Professor Päivi Pajukanta
Department of Human Genetics
David Geffen School of Medicine at
UCLA, Los Angeles, USA

and

Academy Professor Leena Peltonen-Palotie
Department of Medical Genetics
University of Helsinki and
Department of Molecular Medicine
National Public Health Institute
Helsinki, Finland

Reviewed by:

Professor Petri Kovanen
Wihuri Research Institute
Helsinki, Finland

and

Docent Katriina Aalto-Setälä
Department of Medicine
University of Tampere
Tampere, Finland

To be publicly discussed with:

Professor Markku Savolainen
Department of Internal Medicine
University of Oulu
Oulu, Finland

Julkaisija - Utgivare – Publisher**Kansanterveyslaitos (KTL)**

Mannerheimintie 166
00300 Helsinki
p. vaihde 09-47441, telefax 09-47448408

Folkhälsoinstitutet

Mannerheimvägen 166
00300 Helsinki
tel. växel 09-47441, telefax 09-47448408

National Public Health Institute

Mannerheimintie 166
00300 Helsinki, Finland
phone +358-9-47441, telefax +358-9-47448408

Publications of the National Public Health Institute, KTL A16/2004

ISBN 951-740-467-0

ISSN 0359-3584

ISBN 951-740-468-9 (pdf)

ISSN 1458-6290 (pdf)

Helsinki University Biomedical Dissertations No. 61

ISSN 1457-8433

<http://ethesis.helsinki.fi>

Cover: Virve Lilja 2004 "You and me"

Yliopistopaino
Helsinki 2004

Twenty years from now
you will be more disappointed
by the things that you didn't do
than by the ones you did do.
So throw off the bowlines.
Sail away from the safe harbor.
Catch the trade winds in your sails.
Explore. Dream. Discover.

Mark Twain

TABLE OF CONTENTS

LIST OF ORIGINAL PUBLICATIONS	7
ABBREVIATIONS	8
ABSTRACT	9
INTRODUCTION	11
REVIEW OF THE LITERATURE	12
1. OVERVIEW OF LIPOPROTEIN METABOLISM	14
1.2 HDL	14
2. Clinical and metabolic characteristics of familial low HDL-C and FCHL disorders	15
2.1 Low HDL-cholesterol	15
2.2 Familial combined hyperlipidemia	16
2.3 The metabolic syndrome and type 2 DM	17
3. COMPLEX TRAITS AND THE ISOLATED POPULATION OF FINLAND	18
4. STRATEGIES FOR MAPPING GENES UNDERLYING COMPLEX TRAITS	23
4.1 Human Genome Project	24
4.1.1 Comparative genomic approach	26
4.2 Qualitative and quantitative traits	26
4.3 Statistical methods for mapping complex disease genes	27
4.3.1 Parametric linkage analysis	27
4.3.2 Non-parametric linkage analysis: Allele-sharing methods	28
4.3.3 Variance component linkage analysis	29
4.3.4 Association analysis	29
4.3.5 Linkage disequilibrium (LD) analysis	30
4.3.6 Genome-wide association studies	30
4.4 Haplotype blocks and the HapMap project	30
4.5 DNA microarray	31
5. GENETICS OF LOW HDL-C AND FCHL DISORDERS	32
5.1 Candidate genes of low HDL-C	32
5.2 Candidate genes of FCHL	34
5.3 Some important candidate genes of low HDL-C and FCHL	35
5.4 Chromosomal loci identified for low HDL-C	37
5.5 Chromosomal loci identified for FCHL	38
5.6 Chromosome 1q21 region	39
AIMS OF THE PRESENT STUDY	41
MATERIALS AND METHODS	42
1. TABLE OF MATERIALS AND METHODS	42
2. SUBJECTS	43
2.1 Study sample 1: Low HDL-C families	43
2.2. Study sample 2: FCHL families	44
3. LIPID MEASUREMENTS	45

RESULTS	47
1. CANDIDATE GENE STUDY OF FINNISH LOW HDL-C FAMILIES (I)	47
2. GENOME-WIDE SCAN OF FINNISH LOW HDL-C FAMILIES (II)	48
3. FINE MAPPING OF SIX CHROMOSOMAL LOCI IN FINNISH LOW HDL-C AND FCHL FAMILIES (III)	49
4. FINE MAPPING OF THE 1q21 REGION IN FINNISH FCHL FAMILIES (IV)	50
DISCUSSION	54
CONCLUDING REMARKS	57
ACKNOWLEDGEMENTS	61
REFERENCES	64
PREVIOUSLY PUBLISHED IN THIS SERIES BY THE DEPARTMENT OF MOLECULAR MEDICINE	77
ORIGINAL PUBLICATIONS	79

LIST OF ORIGINAL PUBLICATIONS

This thesis is based on the following original publications, referred to in the text by Roman numerals (I-IV):

- I Heidi E. Lilja, Aino Soro, Kati Ylitalo, Ilpo Nuotio, Jorma S.A. Viikari, Veikko Salomaa, Erkki Vartiainen, Marja-Riitta Taskinen, Leena Peltonen, Päivi Pajukanta: A candidate gene study in low HDL-cholesterol provides evidence for the involvement of the APOA2 gene and the APOA1C3A4 gene cluster. *Atherosclerosis* 2002, 164:103-111.
- II Aino Soro*, Päivi Pajukanta*, Heidi E. Lilja, Kati Ylitalo, Tero Hiekkalinna, Markus Perola, Rita M. Cantor, Jorma S.A. Viikari, Marja-Riitta Taskinen, Leena Peltonen: Genome scans provide evidence for low-HDL-C loci on chromosomes 8q23, 16q24.1-24.2, and 20q13.11 in Finnish families. *The American Journal of Human Genetics* 2002, 70:1333-1340.
- III Heidi E. Lilja, Elina Suviolahti, Aino Soro-Paavonen, Tero Hiekkalinna, Aaron Day, Kenneth Lange, Eric Sobel, Marja-Riitta Taskinen, Leena Peltonen, Markus Perola, and Päivi Pajukanta: Locus for quantitative HDL-cholesterol on chromosome 10q in Finnish families with dyslipidemia. *Journal of Lipid Research* 2004, 45:1876-1884.
- IV Päivi Pajukanta, Heidi E. Lilja, Janet S. Sinsheimer, Rita M. Cantor, Aldons J. Lusis, Massimiliano Gentile, Xiaoqun Joyce Duan, Aino Soro-Paavonen, Jussi Naukkarinen, Janna Saarela, Markku Laakso, Christian Ehnholm, Marja-Riitta Taskinen and Leena Peltonen: Familial combined hyperlipidemia is associated with upstream transcription factor 1 (USF1). *Nature Genetics* 2004, 36:371-376.

*These authors contributed equally to this work

Publication II also appears in the thesis of Aino Soro-Paavonen (2004)

The original publications have been reproduced with the permission of the copyright holders.

ABBREVIATIONS

ABCA1	ATP-binding cassette, sub-family A (ABC1), member 1
ApoA1	apolipoprotein AI
ApoA1C3A4	apolipoprotein AI-CIII-AIV
ApoA2	apolipoprotein AII
ApoA4	apolipoprotein AIV
ApoA5	apolipoprotein AV
ApoB	apolipoprotein B
ApoC2	apolipoprotein CII
ApoC3	apolipoprotein CIII
ApoE	apolipoprotein E
ASP	affected sib-pair
bp	base pair
CE	cholesterol esters
CETP	cholesterol ester transfer protein
CHD	coronary heart disease
cM	CentiMorgan
DM	diabetes mellitus
DNA	deoxyribonucleic acid
FCHL	familial combined hyperlipidemia
HDL	high density lipoprotein
HDL-C	high density lipoprotein cholesterol
HL	hepatic lipase
HMG-Co-A	3-Hydroxy-3-methylglutaryl coenzyme A
HRR	haplotype relative risk
IBD	identical by descent
IBS	identical by state
IDL	intermediate density lipoprotein
kb	kilobase
LCAT	lecithin:cholesterol acyltransferase
LD	linkage disequilibrium
LDL	low density lipoprotein
LPL	lipoprotein lipase
LOD	logarithm of odds
PAF-AH	platelet-activating factor acetylhydrolase
PCR	polymerase chain reaction
PL	phospholipids
PLTP	phospholipid transfer protein
PON1	paraoxonase 1
QTL	quantitative trait locus
SNP	single nucleotide polymorphism
SR-BI	scavenger receptor, class B type I
TC	total cholesterol
TDT	transmission/disequilibrium test
TG	triglyceride
type 2 DM	type 2 diabetes mellitus
VLDL	very low density lipoprotein

ABSTRACT

Familial combined hyperlipidemia (FCHL) and serum low HDL-cholesterol (HDL-C) are both highly atherogenic, complex lipid disorders, forming important risk factors for coronary heart disease (CHD), the leading cause of death in Western societies.

FCHL is the most common familial dyslipidemia predisposing to coronary heart disease (CHD) and estimated to be present in about 20% of premature CHD patients. FCHL is characterized by elevated serum total cholesterol (TC), triglycerides (TG), or both, and has many phenotypic features overlapping with the metabolic syndrome and type 2 diabetes mellitus (DM). Low HDL-C is a well-known component trait of both FCHL and type 2 DM. Actually low HDL-C, whether combined with other lipid abnormalities or not, is the most frequent dyslipoproteinemia in patients with premature myocardial infarction. Sharing the phenotypic similarities, these two CHD risk factors are likely to have a close pathophysiologic relationship as well as at least a partially overlapping genetic background.

To analyze the genetic factors underlying low HDL-C and FCHL, we studied multigenerational Finnish families with these lipid disorders and, consequently, premature CHD. We employed several approaches in our analyses. First, we studied candidate genes known to be involved in HDL-C metabolism in 25 low HDL-C families. Among 21 potential candidate genes relevant to lipid metabolism, we found suggestive evidence for linkage with two loci located in the genomic regions of the apolipoprotein A2 (APOA2) gene and the apolipoprotein A1C3A4 (APOA1C3A4) gene cluster.

Second, we performed a genome-wide scan to identify new loci and finally novel genes regulating the metabolic pathways determining serum HDL-C levels. A pooled data analysis of low HDL-C and FCHL study samples provided linkage to 8q23, 16q24.1-24.2 and 20q13.11 with low HDL-C trait. The strongest statistical evidence for linkage was observed on chromosome 8q23 (a lod score of 3.9). Importantly, this locus has earlier been linked to HDL-C levels in Mexican Americans. Further, the region on chromosome 20q has previously been linked to body adiposity, hyperinsulinemia and type 2 DM, suggesting that these traits and HDL-C have a partially shared genetic background.

Next, to further investigate the importance of 8q23, 16q24.1-24.2, and 20q13.11, we analyzed these three low HDL-C loci, as well as two loci for premature CHD on 2q31 and Xq24, and a locus for serum TGs on 10q11, in an extended study material. In this combined study sample of 92 low HDL-C and FCHL families, we fine mapped these six loci and tested their potential involvement especially in quantitative lipid traits. The quantitative HDL-C trait provided strong evidence for linkage to 10q11 (a lod score of 3.3). Furthermore, this region showed evidence of association for a quantitative trait obtained by combining quantitative HDL-C and TGs (p-value 0.0006). The data suggest that the locus on chromosome 10q11, previously linked to TGs, BMI and obesity, influences variation in plasma HDL-C and TG levels in Finnish dyslipidemic families.

The first major locus for FCHL on chromosome 1q21–q23 was identified in a genome-wide scan of Finnish FCHL families in 1998. Since then, this region has been replicated

in FCHL families originating from other populations. Interestingly, the same region has also been linked to type 2 DM in multiple studies. Since FCHL and type 2 DM share several phenotypic features, these data suggest that the same gene may underlie the obtained linkage results. To identify the FCHL gene on 1q21, we sequenced four functionally relevant regional candidate genes, human thioredoxin interacting protein (TXNIP), upstream transcription factor 1 (USF1), retinoid X receptor gamma (RXRG), and APOA2. We also genotyped 60 single nucleotide polymorphisms (SNPs) for 26 genes residing in the linked region. In these analyses, significant evidence for association, linkage and shared haplotypes was found between USF1 and FCHL ($p=0.00002$), especially in males with high TGs ($p=0.0000009$). Furthermore, expression profiles of fat biopsies of FCHL cases differed depending on their carrier status for the associated USF1 haplotype.

In conclusion, USF1 seems to confer susceptibility to high serum lipid levels. Although additional studies are warranted to address the functional differences between different USF1 alleles and their relevance for the FCHL phenotype, our results open novel insights into the genetic background of FCHL, the most common familial lipid disorder predisposing to CHD.

INTRODUCTION

Low serum HDL-C and FCHL represent important risk factors for atherosclerotic vascular disease, and are estimated to explain about 50% of familial dyslipidemias predisposing to premature CHD. Both are typical complex disorders, influenced by several environmental and genetic factors. FCHL is characterized by elevated levels of serum TC and TGs, and also expresses low HDL-C as a component trait. Low HDL-C and the lipid abnormalities of FCHL also overlap with the phenotypic features observed in the metabolic syndrome, a condition characterized by a clustering of risk factors for CHD. The metabolic syndrome significantly increases the risk of death from cardiovascular causes.

When searching for susceptibility genes underlying multifactorial disorders such as low HDL-C and FCHL, the relatively isolated Finnish population is likely to offer some advantages. The Finnish gene pool owes its special characteristics to its relative isolation and unique history of inhabitation involving multiple genetic bottlenecks, which have resulted in a founder effect and increased genetic drift (Norio 2003a; Norio 2003b). Thus, it can be assumed that, in this population, fewer predisposing genes and alleles underlie even complex diseases than in more mixed populations.

The Human Genome Project was completed in April 2003. Human genome sequence data accelerate the search for genes involved in complex traits. The new data freely available for everyone in World Wide Web have also been invaluable for this study by enabling utilization of new methods and strategies in dissection of the genetic background of common dyslipidemias.

The main purpose of this study was to investigate the genetic factors determining low HDL-C and FCHL in multigenerational Finnish families with these lipid disorders and premature CHD. We detected several loci for these lipid abnormalities, and finally identified a novel susceptibility gene for FCHL. This gene, *USF1*, located on 1q21, also offers an interesting candidate for the metabolic syndrome and type 2 DM which have been linked to this chromosomal region in several previous studies.

REVIEW OF THE LITERATURE

1. OVERVIEW OF LIPOPROTEIN METABOLISM

Plasma lipids are transported as water-soluble lipoproteins in the circulation. Lipoproteins consist of a hydrophobic core of cholesterol esters (CE) and TGs, surrounded by surface of more polar components of apolipoproteins, free cholesterol and phospholipids (PL). Plasma lipoproteins are separated into five major classes by ultracentrifugation according to their densities (Table 1) (Gotto et al. 1986).

Table 1. The principal lipoproteins (modified from Ganong 1997).


LIPOPROTEIN	ABBR	DENSITY (g/ml)	COMPOSITION %					PRIMARY APOs	ORIGIN
			Prot	FC	CE	TG	PL		
Chylomicrons	Cm	<0.93	2	2	3	90	3	apoB48, apoC-II	Intestine
Very low density lipoproteins	VLDL	0.94-1.006	8	4	16	55	17	apoB100, apoC-II	Liver, intestine
Intermediate density lipoproteins	IDL	1.007-1.019	10	5	25	40	20	apoB100, apoE	VLDL
Low density lipoproteins	LDL	1.019-1.063	20	7	46	6	21	apoB100	IDL
High density lipoproteins	HDL	1.063-1.210	50	4	16	5	25	apoA-I, apoA-II	Liver, intestine

APOs indicates apolipoproteins; CE, cholesterol esters; FC, free cholesterol; PL, phospholipids; Prot, proteins; TG, triglycerides.

An overview of lipoprotein metabolism is shown in Figure 1. The largest lipoprotein particles, chylomicrons (Cm), are formed in the mucosal cells of the intestine during the absorption of digested exogenous (dietary) lipids and secreted bile in the gut. Cm are cleared from the circulation by the action of lipoprotein lipase (LPL). This enzyme, located on the surface of the endothelium of the capillaries, catalyzes the breakdown of the TGs in the Cm to free fatty acids (FFA) and glycerol, and thus modifies Cm to chylomicron remnants. These cholesterol-rich Cm remnants are taken up by the liver. Delayed remnant removal from plasma by the liver has been suggested to be associated with premature atherosclerosis (Mamo et al. 1998).

Cm provide peripheral tissues with an important source of energy through the β -oxidation of fatty acids. The excess energy of TGs is stored in adipose cells through the reesterification of FFA. During fasting, a portion of these TGs, derived from adipose tissue, re-enter the circulation as FFA and are transported to the liver where they can be used for energy or re-packed into very low density lipoproteins (VLDL). VLDL particles are formed in the liver and contain primarily TGs and CE. These particles are secreted into the circulation where they are then modified by LPL into intermediate density lipoproteins (IDL). IDL can acquire CE from HDL through the mutual exchange of TG by the action of the plasma enzyme cholesterol ester transfer protein (CETP). Some IDL are taken up by the liver and the remaining IDL are further modified into low density lipoproteins (LDL) after losing more TGs by the hydrolytic activity of LPL and hepatic lipase (HL) (Demant et al. 1988).

Figure 1. Overview of the lipoprotein metabolism.



ABCA1 indicates adenosine triphosphate-binding cassette transporter 1; A-I, apolipoprotein A-I; CE, cholesterol esters; CM, chylomicrons; CMR, chylomicron remnants; FC, free cholesterol; FFA, free fatty acids; HDL, high-density lipoprotein; HL, hepatic lipase; IDL, intermediate-density lipoprotein; LDL, low-density lipoprotein; LDL-R, low-density lipoprotein receptor; LRP, low-density lipoprotein receptor-related protein; LPL, lipoprotein lipase; PGN, proteoglycans; PL, phospholipids; SRA, scavenger receptor; SR-B1, scavenger receptor B1; VLDL, very-low-density lipoprotein.

LDL provide cholesterol to the tissues via LDL-receptor mediated uptake, where apoB100 acts as a ligand. Oxidized LDL is also taken up by the scavenger receptor of the macrophages (Brown et al. 1980). Macrophages overloaded with oxidized LDL are called foam cells, a cellular component of atheromatous plaque.

The small intestine and liver additionally synthesize nascent HDL, which returns excess cholesterol from peripheral tissues to the liver (reverse cholesterol transport) for excretion from the body in the bile (Rader 2003). In reverse cholesterol transport, HDL absorb cholesterol from peripheral tissues and importantly, from macrophages/foam cells (Chinetti et al. 2000; Brewer and Santamarina-Fojo 2003). CE of HDL particles are transferred through CETP to TG-rich lipoproteins, namely to IDL and LDL, and, eventually, back to the liver. The increased transfer of CE from HDL to IDL, and from LDL to VLDL, in return for TGs, contribute to the formation of highly atherogenic, small-dense LDL particles (Austin et al. 1990b; Krauss 1994; Packard et al. 2000; Kwiterovich 2002; Deeb et al. 2003).


1.1 HDL

High density lipoproteins, the smallest particles of the five lipoprotein classes present in the circulation, are a heterogeneous population of discoidal and spherical particles of differing size and composition. HDL particles are divided into two major subclasses by ultracentrifugation, HDL3 ($d=1.125\text{-}1.250\text{ g/ml}$) and HDL2 ($d=1.063\text{-}1.0125\text{ g/ml}$). HDL2 particles are about 50% larger and contain more cholesterol and TGs than HDL3 particles (Eisenberg 1984; Gotto et al. 1986).

Two major protein components of HDL-C particles are apolipoprotein-AI (apoA-I) (70%) and apolipoprotein-AII (apoA-II) (20%). ApoA-I is found in all HDL subclasses, whereas several other smaller apolipoproteins, apoA-IV, apoA-V, apoC-I, apoC-II, apoC-III, apoIV, apoD, apoE, and apoJ, are found in mature HDLs (Barter et al. 2003). Additionally, lipid transfer proteins (phospholipid transfer protein, PLTP; CETP), and modifying enzymes (lecithin:cholesterol acyltransferase, LCAT; paraoxonase 1, PON1; platelet-activating factor-acetylhydrolase, PAF-AH) exist in HDL. HDL particles are constantly remodeled by interaction with lipases, lipid transfer proteins, and cell-surface HDL receptors (Figure 2).

Figure 2.

Overview of the metabolic pathways of HDL and reverse cholesterol transport. Dietary cholesterol is not shown. TG indicates triglycerides, CE cholesterol esters, PL phospholipids and FFA free fatty acids.



Many proteins are known to be involved in HDL metabolism and to have a role in the regulation of HDL-C levels (Figure 2). These proteins can be divided into five groups: (1) HDL related apolipoproteins, (2) HDL associated enzymes and transfer proteins, (3)

plasma and cell enzymes that affect HDL, (4) cellular receptors and transporters that interact with HDL, and (5) transcription factors that influence HDL (Wang and Paigen 2002). Multiple mechanisms contribute to the transfer of cellular cholesterol to HDL. These include the ABCA1 transporter, which catalyzes active transfer of cholesterol from the cholesterol-rich cell surface to the HDL particle, and passive (nonspecific) transfer of cholesterol from the cell membrane via intermediates including serum albumin.

2. CLINICAL AND METABOLIC CHARACTERISTICS OF FAMILIAL LOW HDL-C AND FCHL DISORDERS

2.1 Low HDL-cholesterol

A decreased HDL-C level (also called hypoalphalipoproteinemia) is the most common lipoprotein abnormality in patients with premature coronary artery disease (Genest et al. 1992). High levels of HDL-C protect against CHD (Castelli et al. 1977; Gordon et al. 1977; Wilson et al. 1988; Gordon et al. 1989), and low HDL-C is a highly atherogenic disorder. Familial low HDL-C is a complex disorder, where genetic factors have a key role in regulating HDL-C levels (Prenger et al. 1992; Knoblauch et al. 1997). Rare forms of genetic HDL-C deficiency have been described, but the major loci affecting plasma HDL-C levels are still unknown. It has been estimated, that 40 to 60% of the variation in HDL-C levels are caused by genes (Hamsten et al. 1986; Prenger et al. 1992; Bu et al. 1994; Cohen et al. 1994; Guerra et al. 1997; Knoblauch et al. 1997; Inazu et al. 2001).

While genetic factors are major determinants of HDL-C (Hamsten et al. 1986; Cohen et al. 1994), environmental factors also influence HDL-C levels. Environmental factors include unmodifiable contributors such as age and gender. Modifiable lifestyle factors include alcohol consumption (Ellison et al. 2004), physical activity (Durstine et al. 2001), and smoking (Garrison et al. 1978; Criqui et al. 1980). In addition, BMI and TG levels are important determinants of HDL (Anzalone et al. 1995; Devroey et al. 2004), and low HDL-C is common in patients with abdominal obesity, type 2 DM, and the metabolic syndrome.

Decreased HDL-C (<35mg/dl or 0.9 mmol/L) is an independent risk factor for CHD (Schaefer et al. 1994). Therefore, HDL metabolism is an attractive target for drug development in order to elevate HDL levels and decrease the risk for CHD. Currently there are no pharmacological interventions available that specifically raise HDL-C and leave other lipid levels unchanged. However, complex HDL metabolism offers potential opportunities for drug discovery. For example, pharmaceutical approaches to HDL elevation include niacin and the fibric acid derivatives, which both raise HDL-C and have been shown to decrease the risk for CHD (Saefer and Cornell 2000).

The role of HDL as an independent inverse predictor of CHD has been firmly established by numerous epidemiologic studies (Castelli et al. 1986; Assmann et al. 1996). HDL is involved in reverse cholesterol transport, where HDL particles deliver excess cholesterol from the peripheral tissues to the liver or transfer cholesterol into VLDL in a process mediated by CETP (Figure 2). In addition to reverse cholesterol transport, HDL has many other antiatherogenic roles (Table 2) (reviewed in Navab et al. 2001; von Eckardstein et

al. 2001; Assmann and Nofer 2003; Assmann and Gotto 2004). For instance, HDL acts as an anti-inflammatory and antioxidant agent (Table 2).

Table 2.

ANTIATHEROGENIC ROLES OF HDL
Reverse cholesterol transport
Anti-thrombotic properties
Pro-fibrinolytic activity
Antioxidant effect: Prevention of LDL oxidation
Anti-inflammatory effect: Inhibition of endothelial adhesion molecule expression

Interestingly, inflammatory mechanisms are increasingly recognized to play an important role in vascular disease, as inflammatory markers correlate with prognosis in acute and chronic CHD. Recently, HDL was shown to prevent the CRP-induced upregulation of inflammatory adhesion molecules (Wadham et al. 2004). It has been suggested that the antioxidant properties of HDL are carried out by the enzymes associated with this lipoprotein, such as PAF-AH, PON1, and LCAT (reviewed in Navab et al. 2001; Kovanen and Pentikainen 2003).

2.2 Familial combined hyperlipidemia

FCHL, first described in 1973 (Goldstein et al. 1973; Nikkila and Aro 1973; Rose et al. 1973), is one of the most common genetic dyslipidemias. About 10-20% of patients with premature CHD have this lipid disorder (Genest et al. 1992), and the population prevalence of FCHL is approximately 1-2% (Goldstein et al. 1973). In FCHL patients, serum TC, TGs or both are elevated (Goldstein et al. 1973; Nikkila and Aro 1973). FCHL is a common, complex disorder, where several genes, environmental factors and their interactions are involved. The heritability of TC and TGs has been estimated: up to 60% of the variability of these lipid levels is determined by genetic factors (Bucher et al. 1988).

In addition to hypercholesterolemia and hypertriglyceridemia, there are several other important component traits in FCHL, such as other atherogenic lipid abnormalities and features of insulin resistance syndrome. The list of these component traits is shown in Table 3.

Table 3. Component traits in FCHL.

ApoB containing lipoproteins ↑	Sniderman et al. 1980; Brunzell et al. 1983;
Small dense LDL particles ↑	Austin et al. 1990a; Hokanson et al. 1993
Glucose tolerance ↓	Hunt et al. 1989; Aitman et al. 1997;
Abdominal obesity ↑	Bredie et al. 1997a; Vakkilainen et al. 1998;
Insulin ↑	Pihlajamaki et al. 2000
HDL-C ↓	de Graaf and Stalenhoef 1998
Free fatty acids ↑	Castro Cabezas et al. 1993
LPL activity ↓	Babirak et al. 1992

The FCHL phenotype overlaps with several common disorders, such as hyperapobetalipoproteinemia, LDL subclass pattern B (small, dense LDL particles), and most interestingly, the metabolic syndrome and type 2 DM (earlier known as non-insulin-dependent diabetes mellitus, NIDDM) (Ayyobi and Brunzell 2003). Importantly, a potential genetic overlap between FCHL, the metabolic syndrome, and type 2 DM has been suggested (Lewis 2002; Ayyobi and Brunzell 2003), making studies of the genetic background of FCHL relevant for these diseases as well.

2.3 The metabolic syndrome and type 2 DM

The metabolic syndrome is a condition characterized by a clustering of CHD risk factors such as insulin resistance, elevated fasting plasma glucose, abdominal obesity, elevated blood pressure, and atherogenic dyslipidemia, including elevated triglycerides and decreased HDL-C level (Grundy 1999). The National Cholesterol Education Program (NCEP) and the World Health Organization (WHO) recently published definitions for the metabolic syndrome. The criteria for the clinical diagnosis for the metabolic syndrome by the Adult Treatment Panel III (ATPIII) are shown in Table 4.

Table 4. The criteria for the clinical diagnosis for the metabolic syndrome by the Adult Treatment Panel III (ATPIII) (NCEP 2001).

Waist circumference	> 102 cm ♂, >88 cm ♀
TGs	>1.7 mmol/l
HDL-C	<1.03 mmol/l ♂, <1.29 mmol/l ♀
Blood pressure	≥ 130/85 mmHg
Fasting glucose	≥ 6.1 mmol/l

At least three of these criteria have to be fulfilled to diagnose the metabolic syndrome. According to these criteria, the prevalence of the metabolic syndrome is over 20% in US (Park et al. 2003) and about 10 % in Finland (Lakka et al. 2002).

The WHO has also published definitions for the metabolic syndrome. According to the WHO, the metabolic syndrome is defined as insulin resistance or the presence of impaired glucose tolerance or type 2 DM and the presence of at least two of the following: abdominal obesity (waist-hip ratio > 0.90 in males and > 0.85 in females or BMI 30 kg/m²), dyslipidemia (serum TGs ≥ 1.70 mmol/l or HDL-C < 0.9 mmol/l in males and <1.0 mmol/l in females), hypertension (140/90 mmHg), or microalbuminuria (Alberti and Zimmet 1998; WHO 1999).

It has been suggested that in type 2 DM patients, the ATPIII proposal more clearly identifies the burden of CHD associated with the metabolic syndrome as compared to the WHO criteria (Marchesini et al. 2004). The WHO definition, however, seems to have relatively high sensitivity and specificity in predicting diabetes (Laaksonen et al. 2002).

The metabolic syndrome is a proinflammatory- and prothrombotic state, where cardiovascular disease is a major clinical outcome. Two subgroups of patients with the metabolic syndrome are suggested to have a particularly high risk for premature CHD: individuals with type 2 DM or FCHL (Carr and Brunzell 2004). These traits account for

up to 50% of early cardiovascular disease risk in patients with the metabolic syndrome. Furthermore, the metabolic syndrome is a very common disorder. For example, it has been estimated to affect 47 million Americans (NCEP 2001). The metabolic syndrome is also closely related to type 2 DM, which is a serious health problem, reaching epidemic proportions worldwide, especially in developed countries.

Dyslipidemia is a typical feature of type 2 DM (Kannel and McGee 1979), and partly because of the atherogenic lipid profile, CHD is the leading cause of death in type 2 DM (Garcia et al. 1974). The most common lipid abnormalities present in type 2 DM are elevated TGs and decreased HDL-C levels (Howard 1987). The lipid abnormalities present in type 2 DM are the same as in the metabolic syndrome, but the dyslipidemic phenotype in type 2 DM tends to be more severe (Taskinen 2003). In addition to dyslipidemia, type 2 DM is also characterized by abnormalities in carbohydrate metabolism (DeFronzo 1997).

As in FCHL and in the metabolic syndrome, the causes of type 2 DM are also multifactorial, including both genetic and environmental elements (Froguel and Velho 2001). The typical features in the pathogenesis of type 2 DM are reduced beta-cell dysfunction and insulin sensitivity, which, together with an obesity-promoting lifestyle and genetic susceptibility alleles, predispose to the disease.


The incidence and prevalence of type 2 DM is increasing almost epidemically due to ageing of the population structures in developed countries, globally increasing obesity and sedentary lifestyles. It has been estimated that, by the year 2010 the total number of people with diabetes will reach 220 million (Amos et al. 1997). Moreover, the worldwide number of cases with diabetes is predicted to double by the year 2025 from a currently estimated 150 million to 300 million (King et al. 1998; WHO 2003). The greatest number of cases is expected in developing countries such as China and India (King et al. 1998).

3. COMPLEX TRAITS AND THE ISOLATED POPULATION OF FINLAND

Complex traits such as FCHL, CHD, DM, and hypertension are consequences of multiple genetic and environmental factors, and their interactions (Figure 3). Such multifactorial genetic disorders, caused by the interactions of multiple genes and the environment, do not follow the classical Mendelian inheritance patterns and are thus called complex traits. Complicated interactions between modifier and susceptibility genes modulate the phenotype of individuals with diseases. For instance, protective alleles can suppress disease in otherwise susceptible individuals. The modifier genes modulate penetrance, dominance, pleiotropy or expressivity in individuals with Mendelian traits and most likely affect complex traits as well (Nadeau 2003).

Figure 3.

CHD and FCHL are typical complex traits where several susceptibility and modifying genes, environmental factors, and their complicated interactions underlie the disease phenotype.



In multifactorial diseases, several typical confounding factors (Table 5) make the investigation of the genetic background exceptionally challenging (Lander and Botstein 1986; Lander and Schork 1994; Risch 2000). For example, complex traits may result more often from noncoding regulative variants than from coding sequence variants (Mackay 2001; Korstanje and Paigen 2002). Thus, to find causal nucleotide variants underlying complex traits, sequencing of coding regions is not enough. Also interpreting the consequences of noncoding sequence variants is complicated, because the relationship between a promoter or intergenic sequence variation, gene expression level, and a trait phenotype is not as well understood as the relationship between the coding variants and protein function (Glazier et al. 2002).

Table 5.

FACTORS COMPLICATING GENETIC ANALYSES OF COMPLEX TRAITS
Difficulties in diagnosis and classification of the phenotype
Variable expressivity of the phenotype
Late onset of the disease
Quantitative phenotypes
Unknown mode of inheritance
Phenocopies
Unknown or low penetrance
Pleiotrophy
Epistasis
Common disease-predisposing alleles
Genetic heterogeneity
Limited statistical power
Multiple testing
Publication bias

Some efforts can be made to reduce the impact of confounding factors in the background of common complex diseases. For instance, focusing on familial cases with well-defined clinical phenotypes, early onset of the disease and severe phenotype, and exclusion of obvious environmental risk factors should enrich the genetic forms of the disease. Concentration on families originating from population isolates as well as investigation of animal models can further help tackle the genetic background of complex diseases and may help dissect the molecular mechanisms underlying these disorders.


Genetic and environmental heterogeneity are reduced in genetically isolated populations such as Finland (Peltonen et al. 1995), Sardinia (Angius et al. 2001), Costa Rica (Mathews et al. 2004), the Canadian Hutterites (Ober et al. 2001), the Old Order Amish (Cross 1976), the North American Mennonites (Jaworski et al. 1989), the Ashkenazi Jews (Permutt et al. 2001), and the Benkala in Bali (Friedman et al. 1995). When studying individuals originating from population isolates, the impact of genetic involvement is more homogeneous than in mixed populations. Thus, possibilities of identifying contributing genes for common lipid disorders such as FCHL are likely to be increased. It has been suggested that stable populations such as the Saami should be well suited for mapping complex traits, whereas rapidly expanded populations such as the Finns for mapping monogenic diseases (Laan and Paabo 1997; Terwilliger et al. 1998).

Although there have been some inhabitants in Finland since the end of glacial period about 11 000 years ago (Nevanlinna 1972; de la Chapelle 1993), Finland has mainly been inhabited in two waves. Early settlement occurred more than 2000 years ago and populated mainly the coast and the southern part of Finland (Nevanlinna 1972; de la Chapelle 1993). Late settlement took place during the 16th century, from a small south-eastern area to the central, western and finally northern parts of the country, thus inhabiting wilderness and yet non-inhabited parts of Finland (Figure 4). This internal migration movement in the 1500s produced internal isolates and regional subisolates. Because of the persistence of the subisolates, the prevalence of some mutations differs regionally, and local clusterings of some disease alleles exist (Pastinen et al. 2001).

According to another inhabitation theory called the dual-origin hypothesis, two different groups settled Finland: one from east, and the other from south (Eriksson 1973; Norio et al. 1973). Genetic evidence from Y chromosome supports this dual-origin model (Kittles et al. 1998). Despite the opposing inhabitation theories it has been proposed that the current Finnish population is related to other European populations (Sajantila et al. 1995; Lahermo et al. 1996). A recent review article by Norio (Norio 2003b) discusses about the settlement of Finland and summarizes that the Finns are about 10 000 year-old north-Europeans, who also have genes from southern Europe and additionally, some genes as a result of movement in east-west direction.

For geographical, linguistic and cultural reasons, the Finnish population remained isolated for about 2000 years. In addition, multiple bottlenecks (such as wars, diseases, famine years) molded the gene pool. The Finnish population has also grown exponentially from 18th century to date, from 250 000 inhabitants to 5.2 million.

Figure 4. The internal migration movement of the 16th century. (Modified from Peltonen et al. 1999)



The relatively isolated population of Finland has been shown to offer many advantages for genetic research and studies of monogenic traits (Peltonen et al. 1999), and these benefits might also help to dissect complex disorders (Peltonen et al. 2000). In addition to isolation, the founder effect and genetic drift have also had an impact on the population. In the founder effect, a new colony is formed by a very small number of individuals from a larger population and this produces a loss of genetic variation. In genetic drift, changes in the frequencies of alleles in a population occur by chance, rather than because of natural selection. The magnitude of the gene frequency changes due to genetic drift is inversely related to the size of the population; the larger the number of reproducing individuals, the smaller the effects of genetic drift.

Why do harmful alleles exist in a population? Gene selection in the past has probably been influenced by conditions of life in those days, for example famine (Damcott et al. 2003; Chakravarthy and Booth 2004). To ensure survival during periods of famine, some alleles that may have been advantageous in ancient times may have concentrated in population. This heterozygote advantage means that the high rate of occurrence of a condition that is usually very harmful or even lethal in its homozygous form may allow greater survival for heterozygotes for example during famine. Thus heterozygotic carriers of the gene are more likely to survive to adulthood and therefore to pass their genes to the next generation.

The alleles that used to be advantageous in the distant past may today be disadvantageous and cause for instance obesity and predispose to CHD because of the lifestyle of the modern-day humans including physical inactivity and a positive caloric balance. The genes that constitute a risk for developing obesity-related conditions, such as type 2 DM and the metabolic syndrome, when energy is abundant and that on the other hand, conserve energy during famines are called thrifty genes. “Thrifty genotype”, defined as being exceptionally efficient in the intake and/or utilization of food, was initially proposed by Neel (Neel 1962), who argued that certain genotypes were selected into the

human genome because of their selective advantage over the less “thrifty” ones. The FOXC2 gene residing in the 16q region linked to low HDL-C (Pajukanta et al. 2003) has been proposed to work as an anti-thrifty gene in hypertriglyceridemia, obesity, and diet-induced insulin resistance (Cederberg et al. 2001). A high fat diet induces FOXC2 expression in adipocytes, leading to a lean and insulin sensitive phenotype. Thus, increased FOXC2 levels could be protective against type 2 DM (Cederberg et al. 2001).

Due to the founder effect and to isolation, monogenic diseases are less likely have both locus and allelic heterogeneity, and affected individuals tend to share ancestral haplotypes derived from a handful of founders. Thus, strategies using linkage disequilibrium (LD) and a shared haplotype in affected individuals can be applied to restrict the critical DNA region efficiently (Peltonen et al. 1999; Peltonen et al. 2000). In fact, LD has been successfully used in mapping of genes for numerous rare monogenic diseases which belong to the Finnish disease heritage (Peltonen et al. 1999). However, later on it has become evident that monogenic traits are often not as simple as was expected (Nabholz and von Overbeck 2004). Several different mutations in the same or in different loci, with variable phenotypic effects and highly variable associated risks have been identified underlying diseases of simple Mendelian inheritance. For example, the Marfan syndrome has turned out to be fairly heterogeneous, showing both locus and allelic heterogeneity. The Marfan syndrome can be caused by mutations of two different genes and, on the other hand, patients with the same mutation can show a wide degree of phenotypic variability (Collod-Beroud and Boileau 2002).


As in monogenic diseases, there might be fewer disease predisposing alleles also in multifactorial diseases in the relatively isolated population of Finland, compared to populations of more heterogeneous origin (Lander and Botstein 1986; Lander and Schork 1994; Collins 1995; Peltonen et al. 2000). Accordingly, Finnish study samples, consisting of many large pedigrees with multiple affected individuals, have been used in genetic studies of multiple complex diseases (Table 6). These studies have in most cases resulted in the initial positioning of several susceptibility loci, and many of these loci have also been detected in other populations. For example, the first major locus for FCHL in 1q21-23 was detected in Finnish FCHL families (Pajukanta et al. 1998). Since then, this linkage has also been replicated in other, more heterogeneous study samples (Coon et al. 2000; Pei et al. 2000; Allayee et al. 2002).

Table 6. Examples of genome scans for common traits performed previously utilizing Finnish study samples in the National Public Health Institute, Helsinki, Finland

TRAIT	AUTHORS AND YEAR
Multiple sclerosis	Kuokkanen et al. 1997
Schizophrenia	Ekelund et al. 2000; Paunio et al. 2001
Bipolar disorder	Ekholm et al. 2003
FCHL	Pajukanta et al. 1999
Low HDL-C	Soro et al. 2002
Premature CHD	Pajukanta et al. 2000
Hypertension	Perola et al. 2000
Osteoarthritis	Leppavuori et al. 1999
Obesity	Ohman et al. 2000
Autism and Asperger syndrome	Auranen et al. 2002; Ylisaukko-oja et al. 2004

Importantly, in population isolates like the Finns, a high degree of environmental homogeneity is also typical (Figure 5) (Peltonen et al. 2000). For example, life style, diet and culture tend to be more similar among isolated than in more mixed populations. In general, Finns also seems to have a positive attitude towards medical research. In addition, uniformity in physician training and clinical practice yields consistent diagnoses and phenotyping of diseases.

Figure 5. Advantages for genetic research in Finland: Reduced genetic and environmental heterogeneity.



4. STRATEGIES FOR MAPPING GENES UNDERLYING COMPLEX TRAITS

Since complex diseases do not follow classic and simple Mendelian inheritance patterns, the genetic dissection of complex traits is very challenging. Glazier et al. suggested a four-step strategy for gene discovery in studies of complex traits (Figure 6) (Glazier et al. 2002).


In step 1, a whole-genome linkage study is performed. Statistically significant evidence of linkage has to be established in a single study or consistent suggestive evidence has to be found in several independent studies (Lander and Kruglyak 1995; Risch and Merikangas 1996; Dahlman et al. 2002). Because of the nature of the complex traits, the minimal interval for the restricted locus in primary genome screens is usually no less than 10 to 30 cM (containing ~ 100 to 300 genes), a resolution that is sufficiently precise to justify further study (Glazier et al. 2002).

In step 2, the aim is to reduce the size of the critical region. This can be done in humans by utilizing genetic methods such as LD mapping, family-based studies, and case-control studies. These approaches may even reduce the minimal interval to 1 cM.

Identification of the candidate nucleotide variants residing in the region is performed in step 3. Nowadays this can be mainly done by utilizing SNP databases, and burdensome large-scale DNA sequencing analysis of the region is not needed any more. After identification of the SNPs, the relevance of each candidate nucleotide variant and haplotype has to be tested.

In step 4, to get conclusive evidence, associated candidate genes are functionally tested. The gene plays an important role in the trait, if replacement of the variant nucleotide results in changes in phenotype. This proves that a molecular variant is functionally associated with differences in phenotype. Functional tests can be carried out by traditional cell biology methods, such as overexpression studies in transfected cell lines, or by transgenic and gene-targeting technologies. For cellular phenotypes even in vitro functional tests may be appropriate (Glazier et al. 2002).

Figure 6. Overview of the a four-step strategy for gene discovery in studies of complex traits (Glazier et al. 2002).



Although the four-step strategy mentioned above has been commonly utilized previously, the strategy may change in the future, after the International HapMap Project has produced a haplotype map of the human genome. The HapMap will describe the common patterns of human DNA sequence variation and this information might facilitate genome-wide association studies in the search for genes behind complex traits (see 3.2.5 Genome-wide association studies).

4.1 Human Genome Project

In 2001, as a result of international collaboration in the Human Genome Project (HGP), a draft sequence of the human genome became freely available (Lander et al. 2001). At the

same time, the sequence of the human genome was also published by the Celera company (Venter et al. 2001). Since HGP was completed in April 2003, all of the sequence data generated by the International Human Genome Sequencing Consortium has been swiftly deposited in public databases and made freely available to scientists around the world (<http://www.ncbi.nlm.nih.gov/genome/guide/human/>).

Although HGP is essentially completed, the exact number of genes encoded by the genome is still unknown. The latest estimates from gene-prediction programs suggest that there might be about 24 500 or fewer protein-coding genes (Pennisi 2003), much fewer than the previous estimates of around 100 000. This smaller number of human genes has been a surprise, because with around 30 000 genes, the human gene count is only about 30% greater than that of the simple roundworm *C. elegans* with about 20 000 genes (Table 7) (Claverie 2001), and counting genes has been viewed as a way of quantifying genetic complexity. However, people and chimpanzees have roughly 98 percent of their DNA in common; suggesting that the biological consequences due to the genetic differences are much more complicated than was previously speculated (Weissenbach 2004).

Table 7. The comparative genome sizes of human and other organisms sequenced.

Organism	estimated size	estimated gene number
<i>Homo sapiens</i> (human)	3000 million bases	~30,000
<i>Mus musculus</i> (mouse)	3000 million bases	~30,000
<i>Drosophila melanogaster</i> (fruit fly)	180 million bases	13,600
<i>Arabidopsis thaliana</i> (plant)	125 million bases	25,500
<i>Caenorhabditis elegans</i> (roundworm)	97 million bases	19,100
<i>Saccharomyces cerevisiae</i> (yeast)	12 million bases	6300
<i>Escherichia coli</i> (bacteria)	4.7 million bases	3200

The smaller number of human genes than was expected is consistent with the theory that variations in gene regulation and the splicing of gene transcripts produce many distinct functions for one protein in different tissues. It also seems that mutations in the coding sequences of genes are responsible for only a delimited number of the differences in disease susceptibility between individuals. In fact, sequence variants (SNPs) that affect gene splicing and regulation must play an important role in determining disease susceptibility (Peltonen and McKusick 2001). As only a few of the millions of SNPs in the genome have such functional impacts, identifying these SNPs is important. Therefore, one of the goals of HGP is to create SNP maps of the human genome.

It has been estimated that a SNP occurs every 200 to 300 bases along the 3.2-billion-base human genome, making up about 90% of all human genetic variation, and resulting in an estimated 11-15 million SNPs (Kruglyak and Nickerson 2001; Salisbury et al. 2003). SNP maps may help to identify the multiple genes associated with complex diseases, in which a single altered gene may make only a small contribution to the disease and therefore associations are difficult to establish with conventional gene-hunting methods (Chakravarti 2001). However, to understand how SNPs and other genetic variations influence the phenotypes and expose the individuals to diseases will be the major challenge for biomedical researchers in the next decades.

4.1.1 Comparative genomic approach

High-throughput genome sequencing of HGP has created a new era for biomedical research, and now the sequencing of entire genomes (the genomes of more than 160 organisms have been sequenced to date) has set the scene for gene discovery. However, while HGP describes the nucleotide composition of nuclear DNA and the anatomy of the genome, it tells us nothing about the phenotypes encoded in the genotypes. Thus, HGP gives us only the object and the tools to further define the real molecular background of diseases.

The increasing availability of genomic sequences from different species enables comparative genomic approaches (Nobrega and Pennacchio 2003; Pennacchio 2003; Pennacchio and Rubin 2003b), which has proved to be a powerful way for the functional annotation of the human sequence. Comparative genomic strategy also forms a feasible tool in the analysis of cardiovascular-related genes (Cheng and Pennacchio 2003).

A basic hypothesis of comparative genomics is that evolutionarily conserved sequences are functionally important, and thus have remained unchanged through time (Nardone et al. 2004). Based on this hypothesis, focusing on sequences that are highly conserved between different species, for example between human and mouse, can offer powerful tools for the discovery of biologically active gene regulatory elements and facilitate the identification of genes behind complex traits. This strategy was utilized to identify additional functional elements in the vicinity of the APOA1C3A4 gene cluster and, as a result, led to the identification of a novel apolipoprotein member of this gene cluster, APOA5 (Pennacchio and Rubin 2001). The animal model for the trait can also offer potent tools for dissecting the trait through experimental crosses, followed by studies of genetic homology in humans. After the genes are identified in mouse models, their human orthologues can be predicted.

4.2 Qualitative and quantitative traits

Human diseases can represent either a qualitative or quantitative trait. The qualitative trait is a so-called on-off trait, which cannot be measured but rather the disease is either present or not (e.g. albinism). However, quantitative phenotypes often underlie these on-off traits, when they are based on a superimposed cut-off value of the quantitative trait (e.g. hypertension).

Multifactorial diseases and especially their trait components are often quantitative traits showing continued variation, such as blood pressure in hypertension, BMI in obesity, glucose tolerance in diabetes, and plasma lipid levels in lipid disorders. Such measurable characteristics often follow a normal distribution with mean and variance, forming a bell shaped curve. A certain correlation is usually seen in measured values: tendency of the measured values is more similar among relatives than among the general population. This correlation can be either positive or negative, reflecting both genetic and environmental influences.

The locus affecting complex disorders is called the quantitative trait locus (QTL). The QTL is a polymorphic locus which contains alleles that differentially affect the expression

of a continuously distributed phenotypic trait. Generally, it is a genetic marker described by statistical association with quantitative variation in the particular complex trait that is thought to be controlled by the cumulative action of alleles at multiple loci.

Heritability is the fraction of the total phenotypic variance of a quantitative trait caused by genes, estimating the contributions of the different alleles to the variability of a particular quantitative trait. Heritability of a quantitative trait can also be estimated in twin studies, but this parameter does not reveal how many genes are involved or how the different genes interact.

Complex traits such as multiple sclerosis, obesity and CHD can also be analyzed as qualitative traits. In that case, an underlying liability distribution is assumed and, if a threshold of liability is exceeded, an individual becomes affected (polygenic threshold trait). For example in obesity, BMI is either above 30 and a person is obese (affected), or BMI is below 30 and a person is not obese (unaffected). The qualitative trait method has been used successfully in studies of complex disorders, in which families have been identified through an affected individual, otherwise known as a proband. Additional family members are collected to build multigenerational families in which genetic linkage studies can be undertaken.

4.3 Statistical methods for mapping complex disease genes

Several different statistical approaches can be utilized to map the genes involved in the etiology of common complex diseases, including parametric linkage analysis, nonparametric allele-sharing based methods (affected-sib-pair and affected-pedigree-member methods), association analysis, linkage-disequilibrium-based analysis, and polygenic analysis of experimental crosses (Lander and Schork 1994). The choice of the analysis method depends on the study sample available, the trait investigated (e.g. qualitative or quantitative) and prior knowledge of genetic determinants (e.g. positional or functional candidates, model organisms).

4.3.1 Parametric linkage analysis

The basic approach to genetic mapping is linkage analysis, which has been traditionally and successfully applied to simple monogenic traits. It can also be utilized in a linkage analysis of extended families in multifactorial diseases (Curtis et al. 1995). Linkage analysis is a parametric method for testing genetic linkage. In this parametric approach, several parameters need to be estimated: mode of inheritance and frequency of the disease gene, as well as penetrance (meaning the probability of being affected, given the certain genotype). In complex traits, all these parameters are typically unknown, leading to a more or less “wrong” model and, thus, reducing the possibilities and power to detect true linkage signals (Clerget-Darpoux et al. 1986). These problems can be partially avoided in nonparametric methods such as allele sharing methods, in which no specified parameters of the disease locus characteristics are needed.

Linkage analysis determines whether the alleles at the disease locus and at a polymorphic marker locus co-segregate in a pedigree more often than they would if they were located physically far away from each other or even on different chromosomes. In linkage

analysis, it is determined if the recombination fraction (θ) between the two loci differs significantly from $\theta=0.5$, which is expected for unlinked loci. The likelihood-based lod score method (Morton 1955) can be used to analyze linkage within families. This method determines a lod score (Z), which is a log of the odds that two gene loci are linked versus unlinked. In linkage analysis, the overall likelihood of the data for two alternative assumptions is calculated: first, that the two loci are linked with the given recombination fraction (θ), and second that they are not linked. The logarithm to the base 10 of the ratio of these two likelihoods is the lod score (Z) for the “logarithm of the odds”. $Z=\log_{10}$ is the likelihood of data if loci are linked with a given θ .

$$Z(\theta)=\log_{10} \frac{L(\text{linkage})}{L(\text{no linkage})}$$

In the formula above, L is the likelihood function and θ is the recombination fraction. The most likely distance between two loci (e.g. a marker and a disease gene) is the recombination fraction at which the lod score peaks. A lod score of 3.0 corresponds to a p-value of <0.001 (Ott 1991), and is considered to be significant evidence for linkage, whereas a lod score of -2.0 indicates significant evidence for exclusion of linkage when investigating a monogenic disease.

Software programs, such as the LINKAGE package (Lathrop and Lalouel 1984; Lathrop et al. 1984; Lathrop et al. 1986), are commonly used to test linkage in practice. The LINKAGE package includes a series of programs for maximum likelihood estimation of recombination rates, and calculation of two-point and multipoint lod scores, as well as analysis of genetic risks.

4.3.2 Non-parametric linkage analysis: Allele-sharing methods

Non-parametric methods, also called “model-free” methods, partially circumvent some problems associated with linkage analyses. The reason for this is that non-parametric methods are not dependent on the estimates needed in the “model-based” methods, i.e. mode of inheritance, penetrance and gene frequency. In non-parametric methods, all meioses are considered as independent and equally informative for the disease gene. Nonparametric linkage analyses are allele-sharing methods, including affected-sib-pair and affected-pedigree-member methods.

Allele sharing methods are based on allele sharing of affected individuals. Two sibs share 0, 1 or 2 copies of any locus with a 25%-50%-25% distribution expected under Mendelian segregation. Excess allele sharing of a disease-associated locus can be calculated by a chi2 test. Thus, when there is evidence for allele sharing, the affected relatives share the allele(s) more often than is expected by chance. Any two copies of the same allele are called identical by state (IBS). Only if the shared allele is known to be inherited from a common ancestor, is the allele called identical by descent (IBD). Sometimes IBD status cannot be determined unequivocally. IBD methods are more powerful and less prone to errors in allele frequencies than IBS methods, but, especially in late-onset diseases, the parents’ samples are often unavailable, making it impossible to use the IBD methods.

Affected sibpair (ASP) analysis is one of the allele sharing methods. For example, the SIBPAIR program (Kuokkanen et al. 1996) of the ANALYZE package (Terwilliger and Goring 2000) employs this method, in which affected sibpairs are studied to seek distortion from expected allele sharing. Between affected sibpairs, sharing of marker alleles and of phenotypes is more likely when the marker is closely linked to the segregating variation that causes trait variation.

4.3.3 Variance component linkage analysis

It has been suggested that, for quantitative traits such as plasma HDL-C levels, utilizing the full quantitative information of the trait variation is more powerful than arbitrary dichotomizing of the trait into a binary phenotype. There are several methods developed for QTL linkage analysis, such as Haseman-Elston regression (Haseman and Elston 1972) and variance-component methods (Amos 1994; Almasy and Blangero 1998; Blangero et al. 2001). Currently, the variance-component method is one of the most frequently applied approaches in studies attempting to localize QTLs, since, in contrast to regression-based methods, the variance component model can be extended to incorporate covariates, gene-environment interactions and other confounding factors.

The variance-component linkage approach is based on specifying the expected genetic covariances between relative pairs as a function of the identity by descent (IBD) allele sharing relationships at a given QTL (Amos 1994; Almasy and Blangero 1998). Variance component analyses are optimal for large population study samples, where the families are not collected on the basis of some disease. When a sample is non-randomly selected through probands, it might inevitably result in a limited variation for the quantitative trait studied, for instance in serum lipid levels, and thus in an ineffective QTL analysis. Thus, ascertainment correction is generally required. This is done to obtain unbiased parameter estimates when applying variance component methods (Amos and de Andrade 2001).

The variance component approach can be used to analyze large pedigrees, and it is able to accommodate multiple loci simultaneously in a true oligogenic model. The genotype-environment interaction and epistasis can also be examined in a variance component framework. The SOLAR computer package is an example of variance component methods (Almasy and Blangero 1998) that implements general pedigree variance component and IBD estimation methods. SOLAR is able to include non-genetic factors (e.g., age, gender) as covariates. It can also be used to perform multipoint linkage analysis of multiple QTLs.

4.3.4 Association analysis

Association and LD analysis are usually used to analyze candidate genes or to refine the localization of the disease gene, in studies of both monogenic and multifactorial diseases. Association analysis compares the frequency of alleles of the locus between cases and controls. Parental alleles that are not transmitted to the patients can also be used as controls in family-based association analysis, and hence avoid the typical problem of matching control group and cases. The haplotype relative risk (HRR) approach (Falk and Rubinstein 1987; Terwilliger and Ott 1992) and transmission disequilibrium test (TDT) (Spielman et al. 1993) are such family-based methods. In the HRR test, homozygous

parental genotypes also provide some information for the analysis, whereas in the TDT approach they do not. If association is detected, it can mean (1) Direct cause of disease, i.e. the marker is causally implicated in the disease (2) Linkage disequilibrium i.e. the marker is in LD with a susceptibility locus, (3) Natural selection, i.e. there is selection in favor of a certain allele of the marker (4) Population stratification, i.e. the marker allele frequencies are different in the patient and control groups due to population stratification or (5) a False-positive result due to multiple statistical tests.

4.3.5 Linkage disequilibrium (LD) analysis

LD is defined as the non-random association of marker alleles (Slatkin 1994). LD can arise because of several causes, including (1) recent mutation, (2) population founder effects, (3) recent admixture of populations with different allele frequencies and (4) selection in favour of a specific allele (Slatkin 1994). LD mapping is typically used after linkage analysis in mapping monogenic disorders, but genome-wide scans using LD have also been regarded as the method of choice for mapping complex traits (Lander and Schork 1994; Kruglyak 1999; Reich et al. 2001).

4.3.6 Genome-wide association studies

Genome-wide association studies are suggested to be the most efficient way to investigate complex traits (Lander and Schork 1994; Kruglyak 1999; Reich et al. 2001). One reason for this is that association tests are likely to be more powerful than linkage analysis for detecting alleles of modest effect (Risch and Merikangas 1996). New technologies and experimental resources are making whole-genome association studies more feasible, and thus, future approaches to identify genes for complex diseases will include genome-wide association studies. Although whole-genome association studies have good prospects for dissecting the genetics of common diseases, they are currently facing a number of challenges, including problems of multiple testing and study design, definition of intermediate phenotypes and interaction between polymorphisms (Carlson et al. 2004).

4.4 Haplotype blocks and the HapMap project

After fine mapping the region using multiallelic markers and SNPs, haplotypes can be constructed and monitored. The density of markers required depends on the length of LD in the population investigated. Haplotype blocks may help LD mapping by allowing sparser marker maps (Gabriel et al. 2002). The haplotype blocks are sizable regions over which there is little evidence for historical recombination. These haplotype blocks show limited diversity and, in fact, only a few common haplotypes account for over 80% of all haplotypes in the block (Daly et al. 2001; Patil et al. 2001; Gabriel et al. 2002). Therefore, instead of genotyping all SNPs within the haplotype block, only a small fraction, defined as tagging SNPs, are needed to capture most of the haplotype diversity in the population. Such haplotype frameworks, within which only a few common haplotypes are observed, might provide substantial statistical power in association studies of common genetic variation across each region in the future (Gabriel et al. 2002). The constructed haplotype maps of the human genome (www.hapmap.org) may facilitate comprehensive genetic association studies of human disease. However, the optimal SNP density for genome-wide and regional association studies remains to be determined.

The HapMap project (www.hapmap.org) is currently building the haplotype blocks in the human genome. The project is expected to take three years and it involves 9 research groups in 5 countries. The aim is to analyze the block patterns in blood samples taken from people in Japan, Nigeria, and China and from people of northern and western European ancestry in the United States. The HapMap project aims to clarify how the human genome is organized into haplotype blocks. The main purpose is that mapping these haplotypes may form a powerful shortcut to identifying genes for complex traits in the future. The project is largely based on the theory of common variants underlying common disorders (CV/CD hypothesis) (Reich and Lander 2001).

The HapMap is a controversial project among human geneticists because of the questions regarding the block theory. The recent studies (reviewed in Wall and Pritchard 2003) indicate that the structure of LD is “blocklike” in some regions of the genome but not in others, thus some genomic regions fit better to the block theory than others. To summarize, usefulness of the haplotype blocks for future association studies will be uneven across the genome and will depend on the specific patterns of LD in the region of interest (Wall and Pritchard 2003).

With the availability of the human genome sequence and an increasing number of genomic sequences of other species, sequence-based gene discovery will eventually replace the traditional map-based gene discovery (positional cloning) (Peltonen and McKusick 2001). Two future approaches in discovering genetic variations underlying complex disease are a genomic-scale sequence-based approach and a genome-wide haplotype map (utilization of haplotype blocks) (Botstein and Risch 2003). In the sequence-based approach it is assumed that functional SNPs reside in coding regions, splice junctions or promoter regions. It is also expected that nonconservative changes in conserved amino acids are likely to be functional. Thus, this approach can miss functional noncoding SNPs, except when they are evolutionarily conserved. Fewer genotyped SNPs are required as compared to the genome-wide haplotype map approach, and sequence-based strategy also has the potential to detect disease variants with low allele frequencies. Although the sequence-based approach will at least initially focus only on gene regions, which constitute less than 5% of total genomic DNA, it might yield much of the power needed to find the genes underlying complex diseases (Botstein and Risch 2003).

4.5 DNA microarray

Recently, a new technology, called DNA microarray (reviewed in Chittur 2004), has made it possible to monitor the whole genome on a single chip. In a microarray, DNA molecules representing thousands of individual genes are spotted on a microscope slide and thus, researchers can have a picture of the interactions among a considerable number of genes simultaneously. RNA is purified from cells of a particular type and the array is exposed to a labeled sample RNA, hybridized, and the identity/abundance of complementary sequences are determined. Two major application forms for the DNA microarray technology exist: identification of sequence (gene/gene mutation); and determination of expression level (abundance) of genes. Consequently, it is possible to detect which genes are turned on or off in diseased versus healthy human tissues by using microarray technology. The genes that are expressed differently in the two tissues may be involved in causing the disease.

Over the past few years, microarrays have become more widely available to the researchers due to increased commercial availability of custom and generic arrays, the development of robotic equipment that has enabled array printing, and established analysis facilities in research institutions. Currently, microarray technology is a widely employed application for high-throughput comprehensive analysis of gene expression, as well as for genetic analysis and investigations of genomic changes associated with disease.

5. GENETICS OF LOW HDL-C AND FCHL DISORDERS

5.1 Candidate genes of low HDL-C

The inverse association between HDL-C and atherosclerotic cardiovascular disease has long been recognized. Decreased HDL-C, which is the most common dyslipidemia found among patients with premature CHD, is usually combined with other lipid disorders, especially hypertriglyceridemia. Pure isolated familial hypoalphalipoproteinemia (HDL-C < 10th percentile) have been found in 4% of patients with premature CHD as a sole lipid disorder (Genest et al. 1992).

The first observation about the correlation between genetic low HDL-C and precocious atherosclerosis was described by Norum et al (Norum et al. 1982). Paradoxically, monogenic disorders causing low HDL-C do not necessarily correlate with premature CHD. For example, in hypoalphalipoproteinemia caused by mutations in APOA1C3A4, APOA2, LCAT or CETP, no clear association of the genetic defect with CHD have been observed (Assmann et al. 1993; Rader et al. 1993a). Patients with very low HDL-C levels caused by the Apo A-I_{Milano} variant did not have atherosclerotic disease but rather a reduced CHD risk (Franceschini et al. 1980). Moreover, single gene defects are likely to create very low HDL-C levels (HDL-C < 10 mg/dL) without other metabolic problems, while the majority of low HDL-C syndromes, which are caused by multiple genes, exhibit higher HDL-C levels (20-40 mg/dL), and are typically identified in CHD patients (Miller et al. 2003). In addition to CHD, patients having the common form of low HDL-C often exhibit other typical features of the metabolic syndrome as well.

Many genes affecting plasma HDL-C levels have been described (Table 8). However, as in other complex disorders, replications of the identified polymorphisms in other study samples and in different populations have not always been very successful (see Discussion, second paragraph). Interestingly, a recent study suggests that rare alleles with major phenotypic effects significantly influence low HDL-C levels in the general population (Cohen et al. 2004). To determine whether sequence variants collectively contribute to variation in plasma levels of HDL-C, they sequenced three candidate genes causing Mendelian forms of low HDL-C, ABCA1, APOA1 and LCAT, in individuals from a population-based study. In that study, nonsynonymous sequence variants were significantly more common in individuals with low HDL-C than in those with high HDL-C, suggesting that rare DNA sequence variants collectively contribute to variation in HDL-C levels.

Table 8. Genetic defects influencing HDL-C levels.

GENE	LOCUS	SYNDROME	HDL	REPLICATED (+)	REFERENCES
ABCA1	9q31.1	Tangier's disease, Familial HDL deficiency	↓	+	Bodzioch et al. 1999; Brooks- Wilson et al. 1999; Rust et al. 1999
LCAT	16q22.1	Fish-eye disease, LCAT deficiency	↓	+	Funke et al. 1991; Kuivenhoven et al. 1997; Miettinen et al. 1998
APOA1	11q23-q24	HDL deficiency, Familial hypoalphalipoproteinemia, Familial hyperalphalipoproteinemia	↓↑	+	Ordovas et al. 1986; Jeenah et al. 1990; Rader et al. 1993b; Miettinen et al. 1997; Yamakawa- Kobayashi et al. 1999 etc.
APOA2	1q21-q23	Familial apo-A-II deficiency	↔		Deeb et al. 1990
APOC3	11q23.1-q23.2	hypertriglyceridemia	variation		Gagnon et al. 2003
APOA1C3A4 gene cluster	11q23	ApoA-I/C-III deficiency, ApoA-I/C-III/A-IV deficiency	↓		Schaefer et al. 1985; Ordovas et al. 1989
APOA5	11q23	hypertriglyceridemia	↓		Li et al. 2004
LIPC (HL)	15q21-q23	Hepatic lipase deficiency	↑	+	Guerra et al. 1997; Jansen et al. 2002; Su et al. 2002
LPL	8p22	Familial Chylomicronemia syndrome, FCHL, Hyperlipoproteinemia I, Familial LPL deficiency	↓	+	Pimstone et al. 1995; Reymer et al. 1995a
PLTP	20q12-q13.1	No described syndromes in human	↓*		Jiang et al. 2001
CETP	16q21	CETP deficiency	↑		Brown et al. 1989; Zhong et al. 1996; Barzilai et al. 2003
SCARB1 (SR-B1)	12q24.31	No described syndromes in human	↑		Rigotti et al. 1997; Acton et al. 1999

*In mice

5.2 Candidate genes of FCHL

Although FCHL was originally suggested to be a monogenic trait (Goldstein et al. 1973), the inheritance of the FCHL-associated lipid phenotype has later been shown to be complex (Kwiterovich 1993; Aouizerat et al. 1999a). Multiple genes have been shown to play an important role in FCHL and to affect the component lipid traits characteristic of this disorder. Candidate genes of FCHL are shown in Table 9. Juo and colleagues suggested that a common genetic mechanism determines plasma apoB levels and small dense LDL subfraction distribution in FCHL, and a major gene that has pleiotropic effects on these lipid parameters could be the gene underlying FCHL (Juo et al. 1998).

Table 9. Candidate genes of FCHL.

GENE	LOCUS	REPLICATED (+)	REFERENCES
APOA1C3A4- A5 gene cluster	11q13-qter	+	Wojciechowski et al. 1991; Xu et al. 1994; Dallinga-Thie et al. 1996; Deeb et al. 1996; Groenendijk et al. 1999; Groenendijk et al. 2001b; Groenendijk et al. 2001c; Groenendijk et al. 2001d; Eichenbaum-Voline et al. 2004; Mar et al. 2004 etc.
APOA2	1q21-q23		Allayee et al. 2003
APOE	19q13.2		Bredie et al. 1996
LPL	8p22	+	Gagne et al. 1994; Reymer et al. 1995b; de Bruin et al. 1996; Hoffer et al. 1996; Hoffer et al. 1998; Campagna et al. 2002
LIPC (HL)	15q21-q23	+	Gehrisch et al. 1999; Allayee et al. 2000
FABP2	4q28-q31		Pihlajamaki et al. 1997
MNSOD	6q25.3		Allayee et al. 1998; Aouizerat et al. 1999b
LDL-R	19p13.3		Allayee et al. 1998
CETP	16q21		Allayee et al. 1998
LCAT	16q22.1		Allayee et al. 1998; Aouizerat et al. 1999b
USF1	1q22-q23		Pajukanta et al. 2004
TNFRSF1B	1p36.3-p36.2		Geurts et al. 2000

FABP2 indicates the gene encoding Intestinal fatty acid binding protein 2-gene, MNSOD the gene encoding Manganese superoxide dismutase, TNFRSF1B the gene encoding the tumor necrosis factor receptor superfamily, member 1B. APOA1C3A4A5 indicates genes encoding apolipoproteins A-I C-III A-IV and A-V.

Several segregation analysis studies have provided evidence of a gene for elevation of apoB levels (Bredie et al. 1997b). In Dutch FCHL families, evidence for linkage with elevated apoB levels and a locus on chromosome 1p31 was reported (Allayee et al. 2002). A number of variants of the APOA1C3A4 gene cluster (Hayden et al. 1987; Wojciechowski et al. 1991; Groenendijk et al. 2001b; Groenendijk et al. 2001c; Groenendijk et al. 2001d) have also been implicated in the development of FCHL.

Variations in genes affecting the removal rate of TGs from plasma, such as the LPL gene, have been shown to influence the lipid phenotypic expression of FCHL: FCHL individuals with the N291S variant in the LPL gene showed higher plasma lipids and apoB levels compared with affected non-carriers (Campagna et al. 2002). The same variant was associated with decreased HDL-C and increased TG levels in males with FCHL in a study of Reymer et al. (Reymer et al. 1995b), indicating that defective LPL may be one of the factors contributing to the FCHL phenotype. Similarly, two mutations in the LPL gene have been associated with higher plasma TG in FCHL patients in an other study (de Bruin et al. 1996). However, in French Canadian and Finnish subjects with FCHL, the LPL gene mutations showed no association (Marcil et al. 1996; Pajukanta et al. 1997).

5.3 Some important candidate genes of low HDL-C and FCHL

ABCA1

ATP-binding cassette transporter A 1 (ABCA1) is a membrane protein that mediates cholesterol efflux from e.g. macrophages to HDL particles, and thus, takes part in reverse cholesterol transport. The ABCA1 gene resides on chromosome 9q31.1 and is expressed in the liver, the macrophages, and the steroidogenic tissues.

Mutations in the ABCA1 gene cause Tangier disease, a rare autosomal recessive disorder with severe HDL deficiency (Bodzioch et al. 1999; Brooks-Wilson et al. 1999; Rust et al. 1999). Tangier disease has been diagnosed in approximately 40 patients worldwide. In addition to almost complete absence of HDL-C, typical features of Tangier disease include accumulation of cholesterol esters, which leads to enlarged yellow tonsils, hepatosplenomegaly, peripheral neuropathy, and frequently premature CHD (Assmann et al. 1995).

Interestingly, mutations in the ABCA1 gene were also found in French-Canadian patients with familial HDL-C deficiency (Brooks-Wilson et al. 1999). However, a population study of familial HDL deficiency in Finns found no significant association between ABCA1 polymorphism and low HDL-C (Kakko et al. 2003). It is worth noting that familial hypoalphalipoproteinemia patients with ABCA1 mutations have considerably lower HDL-C levels than average CHD patients (Miller et al. 2003). This suggests that HDL-C deficiency caused by ABCA1 variants accounts for only a minority of such cases, and thus, the ABCA1 gene is unlikely to be a major regulator of HDL-C in the general population.

APOAIC3A4 and APOA5

The relationship between variations in the APOAIC3A4 gene cluster and several plasma lipid abnormalities has been well recognized for a long time (reviewed in Groenendijk et al. 2001a). Thus, APOAIC3A4 has been an attractive candidate gene for both FCHL and low HDL-C. Multiple studies predict the importance of the apoAI-CIII-AIV gene cluster as a modifier gene complex in the development of FCHL (Hayden et al. 1987; Wojciechowski et al. 1991; Xu et al. 1994; Dallinga-Thie et al. 1996; Dallinga-Thie et al. 1997; Groenendijk et al. 1999). For example, APOAIC3A4 has been shown to affect LDL-C and apolipoproteins B and C-III (Dallinga-Thie et al. 1996).

The most studied polymorphisms of the APOA1C3A4 gene cluster include three restriction enzyme polymorphisms, XmnI and MspI residing on sites 5' of the APOA1 gene, and the SstI site in the 3' untranslated region of exon 4 of the APOC3 gene (for review see Groenendijk et al. 2001a). The minor alleles of these polymorphisms were associated with elevated plasma cholesterol, TGs, LDL-C, apoB, and apoC-III levels in Dutch FCHL families (Dallinga-Thie et al. 1996). Furthermore, suggestive evidence for linkage between the MspI minor allele and plasma LDL cholesterol levels was detected (Dallinga-Thie et al. 1996). Based on the results obtained in this study, Dallinga-Thie et al. suggested, that the APOA1C3A4 gene cluster is not the primary cause of FCHL, but rather it has a specific modifying effect on plasma triglyceride and LDL cholesterol levels in this lipid disorder (Dallinga-Thie et al. 1996).

In Finnish FCHL families, the MspI polymorphism of the APOA1 gene was associated with total serum cholesterol and apoB levels in spouses, but no evidence of direct involvement of the APOA1C3A4 loci or haplotypes in the expression of FCHL was found (Tahvanainen et al. 1998). Likewise, in French-Canadian subjects with FCHL, no associations were detected between the APOA1C3A4 gene XmnI and SstI polymorphisms (Marcil et al. 1996).

The APOA1C3A4 gene complex has been associated with HDL-C levels in multiple studies (Table 8). Numerous mutations in APOA1 have been described, usually causing decreased HDL-C levels (Ordovas et al. 1986; Yamakawa-Kobayashi et al. 1999), but mutations causing familial hyperalphalipoproteinemia have also been reported (Jeenah et al. 1990; Rader et al. 1993b).

Quite recently, a new member of the APOA1C3A4 gene cluster was proposed: APOA5 (Pennacchio et al. 2001). Since then, growing evidence of the role of APOA5 as an important determinant of plasma TG and cholesterol has been demonstrated, and it seems that APOA5 is also a potent risk factor for cardiovascular disease (Aouizerat et al. 2003; Pennacchio and Rubin 2003a). ApoAV is suggested to reduce plasma triglycerides by inhibiting lipidation of apoB and thus reducing the hepatic VLDL production rate as well as by stimulating LPL-mediated clearance of TG-rich lipoproteins (Schaap et al. 2004).

A strong genetic association has been described between APOA5 polymorphisms and TG levels (Endo et al. 2002; Ribalta et al. 2002; Talmud et al. 2002; Aouizerat et al. 2003; Baum et al. 2003; Martin et al. 2003b; Masana et al. 2003; Seda and Sedova 2003). Over-expression of APOA5 has been shown to lower TGs in mice and APOA5 knockout mice have severe hypertriglyceridemia (van der Vliet et al. 2002). APOA5 is a highly responsive peroxisome proliferator-activated receptor alpha target gene and may act as a major mediator for fibrates in reduction of plasma triglycerides (Vu-Dac et al. 2003).

Linkage and association between variants of the APOA1C3A4A5 gene cluster and FCHL have been demonstrated recently (Eichenbaum-Voline et al. 2004). In more detail, two independent alleles, APOA5c.56G (i.e. S19W) and APOC3c.386G, of the APOA1C3A4A5 gene cluster were overtransmitted in FCHL. In a study of Mar et al., APOA1C3A4A5 showed association with TG levels and LDL particle size in Dutch FCHL families (Mar et al. 2004). The strongest evidence of association in that study was obtained with SNPs in APOA1 (XmnI, MspI) and APOA5 (S19W).

APOA2

The exact function of the second most abundant protein component of HDL-C, ApoA-II, is still unknown. However, polymorphisms in APOA2 have been associated with moderate effects on HDL-C level and/or composition in humans (Scott et al. 1985; Bu et al. 1994; Brousseau et al. 2002; Han et al. 2002). In mice, APOA2 has been associated with HDL-C levels (Hedrick et al. 1993; Mehrabian et al. 1993).

In FCHL, both biochemical and genetic association of plasma apoA-II levels have been suggested: FCHL patients had higher apoA-II levels compared with unaffected relatives, and apoA-II levels showed significant evidence for linkage (LOD=3.1) to a locus on chromosome 1q41 in Dutch FCHL families (Allayee et al. 2003). However, APOA2 is not likely to be a primary genetic determinant of FCHL, as discussed recently by Aouizerat and Kane (Aouizerat and Kane 2003), because the lack of major linkage evidence to the APOA2 region in previous studies (Aouizerat et al. 1999c; van der Kallen et al. 2000). Also in Finnish FCHL families linked to 1q21-23, the peak linkage signal was produced by a marker located about 5 cM from APOA2 (Pajukanta et al. 1998), and association was later detected with variations of the USF1 gene (IV).

TXNIP

A gene for combined hyperlipidemia (Hyplip1) in the mouse has been mapped to a region on mouse chromosome 3 that is potentially orthologous to human chromosome 1q21 (Castellani et al. 1998). The chromosome 1q21 has been identified as an FCHL locus in Finnish, Dutch, German, Chinese and US families (Pajukanta et al. 1998; Coon et al. 2000; Pei et al. 2000; Allayee et al. 2002). The underlying gene, thioredoxin interacting protein gene (TXNIP), which encodes a cytoplasmic protein that inhibits thioredoxin, a regulator of the cellular redox state (Bodnar et al. 2002), thus provided a strong positional candidate for human FCHL (Pajukanta et al. 2001). However, no association between the TXNIP gene and FCHL was found in Finnish FCHL families (IV). Just recently, this negative result has been replicated in Dutch FCHL families (van der Vleuten et al. 2004) as well as in FCHL families participating in the NHLBI Family Heart Study (Coon et al. 2004). Thus, it seems that TXNIP does not play a major role in human FCHL or related traits, and it is unlikely to account for the positive evidence of linkage between FCHL and the 1q21 region.

5.4 Chromosomal loci identified for low HDL-C

Recent family studies have identified six novel loci for this complex disorder (Table 10): Kort et al. found evidence for linkage of the hypoalphalipoproteinemia trait to chromosome 11q23.3 in 105 large Utah pedigrees, approximately 10 cM distal to the apolipoprotein A-I/C-III/A-IV gene cluster (APOA1C3A4) (Kort et al. 2000). In 10 large, randomly ascertained Mexican American families, two novel loci influencing HDL-C levels were localized on chromosomes 8 and 15, using QTL mapping strategy (Almasy et al. 1999). The underlying genes have not yet been identified. A linkage of HDL-cholesterol concentrations to a locus on chromosome 9p has also been demonstrated in Mexican Americans (Arya et al. 2002). The fifth potential locus is the ABCA1 gene on chromosome 9, mutated in the monogenic Tangier disease. ABCA1 can also have allelic variants predisposing to common forms of familial HDL deficiency, as exemplified in

four French-Canadian families with hypoalphalipoproteinemia (Brooks-Wilson et al. 1999). Most recently, a quantitative trait locus influencing plasma HDL-C levels was detected on chromosome 16q in Mexican Americans (Mahaney et al. 2003). This novel QTL locus for HDL-C levels on 16q resides near the loci for LCAT and cholesterol ester transfer protein (CETP). Further evidence of involvement of the 16q locus with a low HDL-C trait has been obtained in a combined data analysis in Finnish and Dutch FCHL families (Pajukanta et al. 2003). In that study, low HDL-C, a component trait of FCHL, was also found to be linked to 9p23.

Table 10. Six potential loci identified for common forms of low HDL-C. Loci marked in bold have been replicated in other study samples.

LOCUS	STUDY SAMPLE	REFERENCES
8q24	Mexican Americans	Almasy et al: 1999
9q31 (ABCA1 gene)	French-Canadian	Brooks-Wilson et al: 1999
11q23	Utah families	Kort et al: 2000
15q21	Mexican Americans	Almasy et al: 1999
16q	Mexican Americans	Mahaney et al. 2003
16q24.1	Finnish and Dutch families	Pajukanta et al. 2003

5.5 Chromosomal loci identified for FCHL

So far, three genome-wide scans have been performed for FCHL in the Finnish (Pajukanta et al. 1999), Dutch (Aouizerat et al. 1999c) and British FCHL families (Naoumova et al. 2003). These scans have identified several chromosomal regions for the FCHL trait or its component traits (Table 11).

Table 11. Loci identified for FCHL in genome-wide scans of FCHL families originating from different populations. Loci that have been detected in several FCHL study samples are indicated in bold.

STUDY SAMPLE	LOCI	REFERENCES
Finnish	1q21 , 2p , 2q, 9p , 10p , 10q, 16q , 20q, 21q	Pajukanta et al. 1998 & 2003, Soro et al. 2002
Dutch	1p, 1q21 , 2p , 9p , 11p , 16q , 19q	Aouizerat et al. 1999, Allayee et al. 2002, Pajukanta et al. 2003
British	6q, 8p, 11p	Naoumova et al. 2003

Some FCHL loci have been replicated in several study samples (loci marked in bold in Table 11). For example, the 1q21 locus has been detected in both Finnish and Dutch FCHL families, as well as in US Caucasians, Chinese and German study samples (Pajukanta et al. 1998; Coon et al. 2000; Pei et al. 2000; Allayee et al. 2002). Another replicated example is the 11p region, which has been detected separately in Dutch and British FCHL families (Aouizerat et al. 1999c; Naoumova et al. 2003). Still, given the expected difficulties in replicating and verifying the results of a genome-wide scan, international collaboration to combine study samples is the utmost importance to accelerate the identification of genes for common cardiovascular traits such as FCHL. This strategy is based on increased statistical power and has been used in previous studies

with other complex traits and it has even led to gene identification in recent studies of asthma (Van Eerdewegh et al. 2002) and inflammatory bowel disease (Hugot et al. 2001; Ogura et al. 2001). This approach was also successfully utilized in a previous pooled data analysis of the Dutch and Finnish FCHL genome scans to identify shared chromosomal regions for FCHL and its component traits (Pajukanta et al. 2003). After unifying the diagnostic criteria and pooling the genotype data, three regions were identified on chromosomes 16q, 2p, and 9p. The 16q24.1 region produced a parametric multipoint LOD score of 3.4 using a parametric location score analysis for the low HDL-C trait. There was also evidence for locus heterogeneity among the families, with about half of them exhibiting linkage to this chromosomal region. Since then, this very same region on 16q has also been linked to HDL-C in an independent study of Mexican Americans (Mahaney et al. 2003).

5.6 Chromosome 1q21 region

The first major locus for FCHL was identified in Finnish FCHL families on chromosome 1q21-23 in 1998 (Pajukanta et al. 1998). This was the first novel locus for FCHL detected since it was described in 1973 (Goldstein et al. 1973). To date, FCHL linkage to 1q21-23 has been replicated in FCHL families originating from many other, more heterogeneous populations (Coon et al. 2000; Pei et al. 2000; Allayee et al. 2002) (Table 12).


Table 12. Traits linked to 1q21-23 (-31)

TRAIT	STUDY SAMPLE	PEAK MARKER/ LOCATION	REFERENCES
FCHL	Finns	D1S104 161,9 cM*	Pajukanta et al. 1998
	U.S. Caucasians	D1S104-D1S1677 161,9-161,5 cM*	Coon et al. 2000
Type 2 DM	Germans	D1S194 165,9 cM*	Pei et al. 2000
	Chinese	D1S194 165,9 cM*	Pei et al. 2000
	Dutch	D1S1665 99,6 cM*	Allayee et al. 2002
	Pimas	D1S1677 161,5 cM*	Hanson et al. 1998
	Utah Caucasians	CRP-APOA2 (165.6–170.8 cM)	Elbein et al. 1999
	French	APOA2-D1S484 158,0-157,5 cM*	Vionnet et al. 2000
	British	D1S2799-D1S452 169,7- 173,2 cM*	Wiltshire et al. 2001
	Amish	near D1S2858 147,2 cM*	Hsueh et al. 2003
	Finns	165 cM	Watanabe et al. 2000
Metabolic syndrome	Chinese	D1S1589 176,25 cM*	Xiang et al. 2004
	Hispanic	D1S518 188,0 cM*	Langefeld et al. 2004
	Caucasians	D1S484 157,5 cM*	Reed et al. 2001
	Western Europeans	150-200 cM	Broeckel et al. 2002
	Plasma FFA	Caucasians APOA2 158,0 cM	Warden et al. 1993

*Estimated sex-averaged genetic location from the p-telomere in centiMorgans according to the deCODE map (<http://www.decode.com/>)

Interestingly, the same markers in the 1q21 region have also been linked to type 2 DM in numerous studies (Hanson et al. 1998; Elbein et al. 1999; Watanabe et al. 2000; Vionnet et al. 2000; Wiltshire et al. 2001; Hsueh et al. 2003), including a Finnish study (Watanabe et al. 2000) (Table 12, Figure 7). Most recently, the very same chromosomal region has been linked to the metabolic syndrome (Langefeld et al. 2004). Therefore, different allelic variants or haplotypes of the same gene may underlie the FCHL, type 2 DM, and the metabolic syndrome linkage results. In addition to overlapping linkage results, these traits share many of their critical metabolic features, further supporting a hypothesis of the partially shared genetic background between these complex disorders.

Figure 7. The critical FCHL area on 1q21-23 and the markers of which have been linked to FCHL, type 2 DM, free fatty acids (FFA), total cholesterol (TC), and the metabolic syndrome (MBS).



AIMS OF THE PRESENT STUDY

The following specific aims were addressed in this study:

- (I) To study the contribution of 21 known candidate genes of HDL-C metabolism for the trait in Finnish low HDL-C families.
- (II) To perform a genome-wide scan to identify novel susceptibility loci for low HDL-C.
- (III) To further investigate the chromosomal loci identified previously in Finnish cardiovascular study samples and to determine the involvement of quantitative lipid traits in Finnish low HDL-C and FCHL families.
- (IV) To fine map and identify the FCHL gene on chromosome 1q21-23.

MATERIALS AND METHODS

1. TABLE OF MATERIALS AND METHODS

The details of the methods used in this study are described in the original publications (I-IV) according to Table 13 below.

Table 13. Materials and methods used in this study.

Material or method	Original publication
Study samples	
Low HDL-C families	I-III
FCHL families	II-IV
Cases and controls	I
Coding as affected subjects	I-IV
DNA extraction	I
PCR and sequencing	I, IV
Statistical analyses	I-IV
Bioinformatics	I-IV
Biochemical analyses	I-III
Genome-wide scan	II
Candidate gene approach	I
Genotyping	I-III
Marker and SNP selection	I-IV
Marker maps	I-IV
Fat biopsies and RNA extraction	IV
Expression array analysis of adipose tissue	IV
Bioinformatics	I-IV
Quantitative real-time PCR analysis	IV
Reporter gene analysis for the transcription efficiency	IV

2. SUBJECTS

The Finnish low HDL-C families were recruited in the Helsinki and Turku University Central Hospitals. The Finnish FCHL families were recruited in the Helsinki, Turku and Kuopio University Central Hospitals. The family collection consisted of three phases. In the first phase, probands were selected from patients undergoing elective coronary angiography or from a registry of patients with confirmed myocardial infarction. In the second phase, probands and all the first-degree relatives were studied to identify low HDL and FCHL families. Finally, in the third phase, all accessible relatives and spouses were contacted and examined, and extensive metabolic and genetic studies were performed. Each subject provided a written informed consent prior to participating in the study. All samples were collected in accordance with the Helsinki declaration, and the ethics committees of the participating centers approved the study designs presented in this thesis.

2.1 Study sample 1: Low HDL-C families (I-III)

Table 14. Diagnostic criteria for low HDL-C probands

INCLUSION CRITERIA
Age 30-60 years
HDL-C < 10th age-sex specific Finnish population percentiles
Premature CHD (angiographically verified >50% stenosis at least in one coronary artery or survived myocardial infarction)
At least three accessible first-degree relatives
At least one first degree relative affected by low HDL-C
EXCLUSION CRITERIA
Diabetes mellitus, type 1 and type 2
BMI >30 kg/m ²
Severe renal or hepatic disease
Hypertriglyceridemia (TG > 2.3 mmol/l for both genders)
Hypercholesterolemia (TC > 6.3 mmol/l ♂, > 6.0 mmol/l ♀)

The following three criteria had to be fulfilled for the myocardial infarction diagnosis: (1) typical clinical symptoms, (2) definite electrocardiographic findings, according to the Minnesota coding (WHO criteria) (Rose et al. 1982), and (3) elevated levels of the creatine-kinase enzyme (CK) and its cardiac isoenzyme, CK-MB.

If the proband fulfilled the criteria mentioned above, families with at least two affected members were included in the study and all the accessible family members were examined. The affected family members were ascertained for low HDL-C, using the Finnish age-sex specific 10th population percentiles (Table 15) derived from the population-based surveys FINRISK and LASERI (Porkka et al. 1994; Vartiainen et al. 1994; Vartiainen et al. 2000).

Table 15. The Finnish age-sex specific 10th population percentiles for HDL-C (mmol/l).

Age (year)	Males	Females
5-11	1.0	1.1
12-14	1.0	1.1
15-17	0.9	1.0
18-20	0.9	1.1
21-24	0.8	1.1
25-29	0.9	1.1
30-34	0.9	1.1
35-39	0.9	1.1
40-44	0.9	1.1
45-49	0.9	1.1
50-54	0.9	1.1
55-59	0.9	1.1
60-	0.9	1.1

2.2 Study sample 2: Finnish FCHL families (II-IV)

Table 16. Diagnostic criteria for FCHL probands

INCLUSION CRITERIA
Age 30-55 years ♂, <60 years ♀
TC and/or TGs >age-sex specific Finnish 90 th percentiles
Premature CHD (angiographically verified >50% stenosis in at least one coronary artery or survived myocardial infarction)
At least one first degree relative affected by FCHL (high TC and/or high TGs) ¹
EXCLUSION CRITERIA
Diabetes mellitus, type 1 ²
Severe renal or hepatic disease
Hypothyroidism
Familial hypercholesterolemia ³

¹If the proband had only one elevated lipid trait, a first-degree relative had to have the combined phenotype.

²Type 2 DM and obesity were not exclusion criteria in the Finnish FCHL families.

³Familial hypercholesterolemia was excluded by determining the LDL-receptor status of the proband by the lymphocyte culture method (Cuthbert et al. 1986).

If the criteria mentioned above were fulfilled, families with at least two affected members were included in the study, and all the accessible family members were examined. FCHL and TGs traits were analyzed. For the FCHL trait, family members were scored as affected if they had the combined hyperlipidemia phenotype or if they had high TC or high TGs using the Finnish age-sex specific 90th percentiles (Table 17). These ascertainment criteria are fully comparable with the original FCHL criteria (Goldstein et al. 1973). For dichotomized analysis of TGs, family members with TG levels \geq 90th Finnish age-sex specific population percentile were coded as affected, and family members with < 90th percentile as unknown.

Table 17.

The 90th age-sex specific population percentiles for total cholesterol (TC), triglycerides (TGs) and apolipoprotein B (apoB) in the Finnish population.

	TC (mmol/l)	TC (mmol/l)	TGs (mmol/l)	TGs (mmol/l)	ApoB (mg/dl)	ApoB (mg/dl)
Age (y)	Men	Women	Men	Women	Men	Women
5-11	5.9	6.2	1.6	1.4	100	108
12-14	6.0	5.8	1.7	1.6	104	103
15-17	5.3	5.6	1.9	1.5	92	101
18-20	5.9	6.0	2.0	1.7	111	111
21-24	5.7	5.9	2.3	1.6	109	115
25-29	6.2	6.1	2.4	1.7	107	102
30-34	6.6	6.2	2.7	1.7	114	106
35-39	7.0	6.4	2.9	1.8	120	109
40-44	7.2	6.6	3.2	1.8	124	112
45-49	7.4	6.8	3.4	1.9	127	115
50-54	7.5	7.1	3.5	2.1	129	119
55-59	7.5	7.3	3.5	2.3	128	122
60->	7.4	7.6	3.5	2.5	127	125

Worth noting is the fact that type 2 DM and obesity were not exclusion criteria for the probands of the families with FCHL. Because of this, some patients fulfilling the criteria of the metabolic syndrome also exist in the study sample.

3. LIPID MEASUREMENTS

All blood samples for the measurement of serum lipids and for DNA isolation were obtained after an overnight fast. Any lipid lowering medication was interrupted for four weeks before the blood samples were taken. All lipid measurements for low HDL and FCHL study subjects were performed according to the same protocols, and moreover, in the same research laboratory of the Division of Cardiology, Department of Medicine, Helsinki University Central Hospital. However, lipid measurements of 6 FCHL families recruited in Kuopio Central Hospital were performed in Kuopio. Serum TC and TGs were measured with an automated Cobas Mira analyser (Hoffman-La Roche, Basle, Switzerland) by enzymatic methods (Hoffman-La Roche kits 0722138 and 0715166, respectively). LDL was calculated using the Friedewald formula (Friedewald et al. 1972):

$$\text{LDL-C} = \text{TC} - \text{HDL-C} - (\text{TG}/2.2)$$

(Valid if TG < 4.5 mmol/L, no apo E2/2 genotype or phenotype, no familial dysbetalipoproteinemia)

Serum HDL-C was quantified by phosphotungstic acid/magnesium chloride precipitation procedures (Hoffman-La Roche kit 0720674). HDL2 and HDL3 were separated by sequential flotation in an ultracentrifuge (Taskinen et al. 1988). Concentrations of Apo-AI, Apo-AII and apoB were measured by immunoturbidimetric methods with commercial kits (Boehringer-Mannheim, Mannheim, Germany). Plasma glucose concentrations were analyzed by the glucose dehydrogenase method (Precision-G Blood Glucose Testing

System, Medisense, Abbott, Illinois, USA). Serum-free insulin concentrations were measured by radioimmunoassay (Phadeseph INSULIN RIA, Pharmacia & Upjohn, Uppsala, Sweden). The homeostasis model assessment for insulin resistance (HOMA IR) was calculated from the fasting plasma glucose and serum insulin concentrations as follows: $\text{fasting insulin } (\mu\text{U/ml}) \times \text{fasting glucose (mmol/l)} / 22.5$ (Matthews et al. 1985; Haffner et al. 1997).

RESULTS

1. CANDIDATE GENE STUDY OF FINNISH LOW HDL-C FAMILIES (I)

The specific aim was to identify and characterize candidate genes affecting the low HDL-C trait in the Finnish low HDL-C families. We analyzed 21 potential candidate genes (Table 18) with 48 flanking markers in 21 low HDL-C families. An additional 4 families were genotyped with the markers for the APOA2 and APOA1C3A4 gene cluster, which showed some evidence of linkage.

Table 18. Genes studied in the candidate gene project of low HDL-C.


CANDIDATE GENE	ABBREVIATION	LOCATION	MARKERS
Apolipoprotein A-I/ C-III/ A-IV gene cluster	APOA1C3A4	11q23	D11S1347, Xmn1, Msp1, Sst1, D11S939
Apolipoprotein A-II	APOA2	1q21-q23	D1S2635, APOA2, D1S2844, D1S104
Apolipoprotein C-II	APOC2	19q13.2	D19S178, APOC2, D19S219
Cholesterol ester transfer protein	CETP	16q13	D16S3057, D16S3143, D16S514
Hepatic lipase	HL, LIPC	15q21-q22	D15S117, LIPC, D15S98
Lecithin:cholesterol acyltransferase	LCAT	16q22.1	D16S3031, D16S496, D16S3139
Phospholipid transfer protein	PLTP	20q12-q13.1	PPGPR, PPGB.PCR
Peroxisome proliferator activated receptor α	PPAR α	22q12-q13.1	D22S928, D22S1160
Peroxisome proliferator activated receptor γ	PPAR γ	3p25	D3S1259, D3S1585, D3S1286
Macrophage scavenger receptor	MSR	8p22	D8S1731, D8S261
Lipoprotein lipase	LPL	8p22	D8S282
HDL-binding protein	HDLBP	2q37	D2S125, D2S395, D2S338
Alpha-1-antitrypsin	A1AT	14q32.1	D14S265, D14S1142
Selectins P, L and E	SELP, SELL, SELE	1q23-25	D1S196, SELP, D1S210
ATP-binding cassette transporter A 1	ABCA1	9q31	D9S277, D9S306
Endothelial lipase	LIPG	18q21.31-q21.33	D18S460, D18S118, D18S1145
Scavenger receptor B 1	SR-B1	12q24.31-q24.32	D12S340, D12S1714, D12S97

In this candidate gene project, we found suggestive evidence for linkage with two loci: Marker D1S2844, located in the APOA2 region, yielded a LOD score of 2.14 and marker D11S939 flanking the APOA1C3A4 gene cluster produced a LOD score of 1.69. We also identified potential shared haplotypes in these two regions in a subset of low HDL families. These results suggest that APOA2 and APOA1C3A4 genes or genes in their immediate vicinity are likely to have an influence on HDL levels. It is conceivable that the obtained linkage result to APOA2 locus on chromosome 1 may as well reflect the presence of the near-by located USF1 gene linked to FCHL in Finnish FCHL families (IV).

2. GENOME-WIDE SCAN IN FINNISH LOW HDL-C FAMILIES (II)

The specific aim of this study was to identify new loci and finally novel genes regulating the metabolic pathways determining serum HDL levels in the Finnish population. We performed a genome-wide scan in three stages (Figure 8). At stage 1, 358 microsatellite markers were genotyped in 20 low HDL-C families. At stage 2, 67 additional markers were genotyped for the regions showing lod scores > 1.0 in stage 1 (15 loci) and 5 low HDL-C families were added to the study sample. The fine mapping provided further support for linkage to the low HDL trait on chromosomes 2p ($Z=2.1$, $\theta=0.10$), 3p ($Z=2.1$, $\theta=0.06$), 8q23 ($Z=2.3$, $\theta=0.10$), 16q24.1-24.2 ($Z=2.2$, $\theta=0.08$), and 20q13.11 ($Z=1.4$, $\theta=0.16$). The significance of these initial findings for low HDL-C was further addressed by analyzing markers in these regions in an independent study sample of 29 Finnish families with FCHL (stage 3), because low HDL-C forms one of the component traits of FCHL. This is also reflected by the fact that there were 64 individuals affected with low HDL-C in these FCHL families.

Figure 8. Overview of the strategy used in the genome-wide scan (study II).



The pooled data analysis of low HDL and FCHL study samples, using low HDL as a tested trait, provided further support for linkage in three of these regions on 8q23, 16q24.1-24.2 and 20q13.11 (Table 19). The highest statistical evidence for linkage was observed on chromosome 8q23 (a lod score of 3.9). Importantly, this locus has earlier been linked to HDL-C in Mexican Americans (Almasy et al. 1999). Further, the region on chromosome 20q has previously been linked to body adiposity, hyperinsulinemia and type 2 DM (Bowden et al. 1997; Ji et al. 1997; Zouali et al. 1997; Ghosh et al. 1999; Lee et al.

1999; Mohlke et al. 2001), suggesting a partially shared genetic background of these traits and HDL-C.

Table 19. The most significant results of the low HDL-C genome-wide search for low HDL-C trait obtained by analyzing the combined study sample of low HDL-C and FCHL families or by analyzing both family sets alone.

LOCUS*	MARKER	STUDY SAMPLE	LOD SCORE
2p25.1	D2S423	FCHL	3.4
2p13.2	D2S1394	low HDL-C	2.1
3p26.1	D3S1304	low HDL-C	2.1
8q23.1	D8S1132	combined	4.7
16q23.3	D16S3091	combined	2.2
20q13.32	D20S171	combined	1.9

* According to the July 2003 human reference sequence of UCSC Human Genome Browser Gateway (<http://www.genome.ucsc.edu/>)

3. FINE MAPPING OF SIX CARDIOVASCULAR LOCI IN FINNISH LOW HDL-C AND FCHL FAMILIES (III)

The aim of this study was to further investigate the significance of six cardiovascular loci identified previously in the genome wide scans of the Finnish study samples in an extended study material consisting of 92 low HDL and FCHL families. The analyzed loci were as follows: low HDL-C loci on 8q23, 16q24.1-24.2, and 20q13.11 (Soro et al. 2002), loci for premature CHD on 2q31 and Xq24 (Pajukanta et al. 2000) and a locus for serum TGs on 10q11 (Pajukanta et al. 1999) (Table 20). We fine mapped these loci and tested their involvement, especially in quantitative lipid traits in a combined study sample of 92 low HDL-C and FCHL families. A total of 67 microsatellite markers were genotyped for these six regions in 1109 individuals (Figure 9).


Table 20. Previous findings from three different genome-wide scans performed in Finnish families¹ or sib-pairs².

STUDY SAMPLE	TRAIT	LOCUS	AUTHORS AND YEAR
FCHL ¹	FCHL	1q21-q23	Pajukanta et al. 1998
	high TGs	10q11	Pajukanta et al. 1999
Low HDL-C ¹	low HDL-C	8q23.1	Soro et al. 2002
		16q23.3	
		20q13.32	
Premature CHD ²	CHD	2q31 Xq24	Pajukanta et al. 2000

TG and HDL-C are highly inversely correlated traits; probably exhibiting at least a partially shared genetic background. To better address this, we analyzed the data of the most promising region of this study (10q11) with a novel QTL association program, which is able to take into account these two QTL traits, TGs and HDL-C, at the same time. In these analyses, we detected evidence for association for quantitative TGs with D10S546, about 7 cM from D10S1772 (p-value of 0.0006). For HDL-C, the association analysis produced a p-value of 0.02 with marker D10S1790, located in the vicinity of

D10S546 (0.0 cM, 889 kb). Furthermore, a combined analysis of quantitative HDL-C and TGs produced a p-value of 0.008 again with marker D10S546 in the association analysis.

Figure 9. Combined study sample of dyslipidemic families.



To summarize the results, we found strong evidence for linkage of the chromosome 10q11 locus with the quantitative HDL-C trait in a combined data analysis, using a variance-component approach (a lod score of 3.3). Furthermore, this region, previously linked to TGs, BMI and obesity (Hager et al. 1998; Pajukanta et al. 1999; Hinney et al. 2000; Ohman et al. 2000; Price et al. 2001), provided evidence of association for both quantitative HDL-C and TG traits. The lowest p-value of 0.0006 was obtained with quantitative TG trait for the marker D10S546. In conclusion, the data suggest that a locus on chromosome 10q11 influences the variation in plasma HDL-C and TG levels in Finnish dyslipidemic families.

4. FINEMAPPING OF THE 1q21 REGION IN FCHL FAMILIES (IV)

The chromosome 1q21 was originally linked to FCHL in Finnish families (Pajukanta et al. 1998). Since then the finding has also been replicated in more heterogeneous populations (Coon et al. 2000; Pei et al. 2000; Allayee et al. 2002). Interestingly, the same region has also been linked to type 2 DM and the metabolic syndrome (Hanson et al. 1998; Elbein et al. 1999; Watanabe et al. 2000; Vionnet et al. 2000; Wiltshire et al. 2001; Hsueh et al. 2003; Langefeld et al. 2004; Xiang et al. 2004). Since FCHL and type 2 DM share several phenotypic features, allelic variants of the same gene may underlie the linkage results obtained. The aim of this study was to identify the underlying FCHL gene on 1q21.

First, a strong positional candidate for human FCHL, the thioredoxin interacting protein (TXNIP) (Pajukanta et al. 2001), was sequenced to identify all possible variants of the gene. All identified 20 SNPs were rare (allele frequencies < 10%) and produced no changes in amino acids. Four TXNIP SNPs was analyzed and, although some evidence for linkage was detected, neither evidence for association or LD, nor evidence for shared haplotypes was found. Thus, the human homolog of the Hyplip1 gene (Castellani et al.

1998), the TXNIP gene, is unlikely to be the causative gene in FCHL, because no statistical evidence emerged in extensive statistical analyses of the identified TXNIP variants or their haplotypes. Furthermore, the human Hyplip1 gene is located about 15 Mb from the peak linkage marker D1S104 in humans, making it a less likely candidate.


Second, three relevant regional candidate genes were sequenced, the upstream transcription factor 1 (USF1), the retinoid X receptor gamma (RXRG), and APOA2. The SNPs that were identified by sequencing the RXRG and APOA2 genes did not result in missense or nonsense variants, and no evidence for association was observed between the SNPs or their haplotypes and FCHL or TGs. Likewise, none of the SNPs that were identified by sequencing of the USF1 gene resulted in amino acid changes. However, two SNPs, usf1s1 (exon 11) and usf1s2 (intron 7), provided evidence for linkage, producing lod scores of 3.5 and 2.0 for FCHL, and 3.7 and 2.0 for TGs. Combined analysis of these two USF1 SNPs produced some evidence for association for both FCHL ($p=0.005$) and the TG trait ($p=0.008$) using the gamete competition test. Usf1s1 and usf1s2 also suggested an association in TG-affected men, and combined analyses of these two SNPs resulted in p -values between 0.02–0.003. Since these SNPs yielded such promising results with the 42 FCHL families genotyped so far, all 60 FCHL families (721 individuals) were genotyped for usf1s1 and usf1s2.

In the analyses of the two USF1 SNPs both the FCHL and TG traits showed evidence for linkage and association ($p=0.00002$ and 0.00006). The strongest evidence for association was seen in TG-affected males ($p=0.0000009$), suggesting some sex-specific effect especially concerning the TG trait. Gender-specific differences are typical for complex traits and both genetic and environmental factors are likely to contribute to these differences. Male sex is a well-known risk factor for CHD, and interestingly, recent studies also suggest gender-specific differences in the dyslipidemic phenotypes in FCHL (Soro et al. 2003).

As mentioned above, the analysis of the two USF1 SNPs, usf1s1 and usf1s2 (Figure 10) provided strong evidence for association. The segregating haplotype in all these analyses was 1-1 (1 indicates the common allele). A total of 15 SNPs were genotyped for the USF1 region. Evidence for association was observed with several SNPs. In the TG-affected males the association extended to the adjacent F11R gene over a 46 kb region, whereas the haplotype analyses in all FCHL- and TG-affected family members restricted the associated region to 1239 bp within the USF1 gene. In addition, haplotype analysis revealed that the 1-1 haplotype of the usf1s1-2 was transmitted significantly more to the TG-affected individuals ($P=0.00003$), especially in TG-affected males ($P=0.0000009$).

In TG-affected men, 6 SNPs in the F11R-USF1 region showed evidence for association, and 5 of these were in strong LD. F11R, also known as JAM1, Junctional adhesion molecule 1, plays an important role in the regulation of tight junction assembly in epithelia. In addition, F11R is involved in the pathogenesis of viral infections and it serves as an integrin ligand. However, the extent of LD in FCHL alleles of both sexes covers only the USF1 gene and the known functions of F11R make it a less likely candidate for FCHL as compared with USF1.

Figure 10. Upstream transcription factor 1 protein is encoded by the 6.74 kb USF1 gene, which maps on chromosome 1q22-q23. Two associated USF1 SNPs are shown in this figure: usf1s1 is located in exon 11 and usf1s2 in intron 7.



To summarize the genetic data, evidence for association, linkage and shared haplotypes was found between USF1 and FCHL. The evidence for association was strongest in TG-affected men. In all the affected FCHL family members, evidence for association ($P=0.00002$) and LD was restricted to the USF1 gene. However, in the TG-affected men, evidence for association ($P=0.0000009$) and LD extends at least 46 kb within the F11R-USF1 region.

Interestingly, preliminary functional data obtained with microarray analysis supported the genetic data: the USF1 risk haplotype had an effect on the expression profiles of fat biopsies. Expression profiles of fat biopsies from affected FCHL family members carrying the 1-1 susceptibility haplotype of USF1 revealed expression differences when compared to affected FCHL family members without this haplotype and homozygous for the putative protective haplotype 2-2. Applying highly stringent criteria, 25 genes were identified as upregulated and 73 genes as downregulated in the risk haplotype carriers. Genes with expression differences included APOE, PLTP, MSP1 and ALOX5. However, no differences in the steady state USF1 expression levels could be observed between FCHL patients ascertained for the risk haplotype, suggesting that the potential functional significance of the associated USF1 allele is not delivered via a direct effect on transcription of this gene in adipose tissue.

To lend biological relevance to the array findings, the lists of differentially expressed genes were examined for any over-representation of functional classes. Only three classes were significantly over-represented among the upregulated genes in the risk haplotype carriers. These were primarily genes involved in fat metabolism. Interestingly, a prominent downregulation of immune-response genes was also observed.

A novel putative Cis-acting regulatory element was identified in intron 7 of USF1, in the immediate vicinity of the USF1 risk haplotype. The genomic sequence flanking the risk haplotype was investigated for potential functional domains, and a 60-bp sequence element present in 91 human genes was identified. The SNP usf1s2, forming part of the risk haplotype, resides adjacent to a 306-bp Alu repeat. Two parts of this Alu repeat show sequence similarity with the mouse B1 repeat. When blasted against the mouse sequence databases, these two parts of the Alu sequence identify numerous mouse ESTs, due to the B1 element located in the untranslated region of the mouse mRNA. A total of 91 human genes, including USF1, have this 60-bp part of Alu located either on the coding strand or

on the opposite strand. The 60-bp part is highly conserved, and found from human to worm but not in *Drosophila melanogaster* or in *Saccharomyces cerevisiae*.

Interestingly, the region in intron 7 of USF1 containing the 60-bp conserved sequence and the usf1s2 SNP showed transcription efficiency. A SEAP reporter assay demonstrated that this 60-bp element has an effect on transcription in vitro in the forward orientation. The reverse orientation resulted in transcription efficiency comparable to that of the negative control, implying a Cis-acting regulator rather than a direction-independent enhancer element. These data suggest a putative regulatory element in the immediate vicinity of the USF1 risk haplotype.

DISCUSSION

The aim of this thesis was to identify disease loci and to search for susceptibility genes for two common atherosclerosis related lipid disorders, low HDL-C and FCHL. First, we detected linkage for low HDL-C near two candidate genes, APOA2 and the APOA1C3A4 gene cluster (I). Second, in a genome-wide scan, three novel loci for low HDL-C were identified, on 8q23, 16q24.1-24.2, and 20q13.11 (II). Third, fine mapping of the previous loci provided more evidence for the importance of the 10q11 region: an association was found between HDL-C and TG levels and 10q11 (III). Fourth, a first susceptibility gene for FCHL, USF1, was identified on 1q21 (IV).

Multiple candidate genes have been suggested to affect plasma HDL-C levels and genes related to FCHL (Tables 8 and 9) and several loci for these lipid traits have been detected (Tables 10 and 11). Typically in complex disorders, the first identification of a genetic polymorphism to be associated with a disease does not automatically lead to replication of the original findings in other study samples and in different populations. The problem in replicating the results may be due to several factors, including phenotypic heterogeneity, inadequate statistical approaches, an insufficient study sample size, or a type 1 error. The failure to replicate a previous positive linkage finding may also be explained by different ethnicity, if the original linked variation is population specific. In future, new statistical methods are needed to better model the complexity caused by several interacting genes and environmental factors.

Replication of the original finding in other study samples and in different populations provides further evidence for the impact of the gene or locus to the trait. Interestingly, some of the results of this thesis study were replications of the previous results or have since then been replicated. Before the detection of linkage of a low HDL-C trait to the APOA2 locus (I), this chromosomal region on 1q21-q23 was linked to FCHL in Finnish (Pajukanta et al. 1998) as well as in some other FCHL families (Coon et al. 2000; Pei et al. 2000). In addition to Finnish low HDL-C families (I), evidence for linkage to near the APOA1C3A4 gene cluster has also been demonstrated in 105 large Utah pedigrees with hypoalphalipoproteinemia (Kort et al. 2000).

Involvement of 8q23 (II) in the regulation of serum HDL levels was earlier shown in randomly ascertained Mexican-American families (Almasy et al. 1999). Further evidence of involvement of a locus on 16q in the regulation of plasma HDL-C levels (II) has been obtained in Mexican Americans (Mahaney et al. 2003) and recently in a combined data analysis of Finnish and Dutch FCHL families (Pajukanta et al. 2003). The 20q13.32 region, which was linked to TGs and HDL-C (II, III), has shown evidence for linkage to type 2 DM and obesity in several studies and in different populations (Bowden et al. 1997; Ji et al. 1997; Zouali et al. 1997; Ghosh et al. 1999; Lee et al. 1999; Ghosh et al. 2000; Mohlke et al. 2001).

The clustering of linkage peaks for component traits of the metabolic syndrome also occur in other chromosomal regions. For example, linkage between 10q11 (III) and obesity has been detected (Hager et al. 1998; Hinney et al. 2000; Price et al. 2001). Likewise, 1q21-q23 is not only linked to FCHL in multiple studies, but also to type 2 DM and the metabolic syndrome (Table 12) (Pajukanta et al. 1998)(Hanson et al. 1998; Elbein

et al. 1999; Coon et al. 2000; Pei et al. 2000; Watanabe et al. 2000; Vionnet et al. 2000; Wiltshire et al. 2001; Allayee et al. 2002; Hsueh et al. 2003; Langefeld et al. 2004; Xiang et al. 2004). Taken together, these findings suggest that one or more genes in this particular chromosomal region may predispose to FCHL, type 2 DM, and the metabolic syndrome, and thus, also to premature CHD. The clustering of linkages in the same chromosomal area is consistent with the fact that many of the critical metabolic features of FCHL (e.g. hypertriglyceridemia and insulin resistance) also represent trait components of type 2 DM and the metabolic syndrome.

The goal in disease gene mapping is to identify new disease genes in order to understand the underlying biological mechanisms and to offer tools for the development of molecular diagnostics, and preventive medicine, as well as drug and possible gene therapy. In a study of FCHL, we have now reached the first step by characterizing a putative susceptibility gene underlying this lipid disorder (IV). Currently we cannot positively confirm one single associated causative variant in the Finnish FCHL families, but rather we identified several associated SNPs in tight LD and a common SNP haplotype, segregating predominantly to affected family members.

FCHL is a typical complex trait, in which several genes, environmental factors and their interactions underlie the disease phenotype. Variants involved in complex disease are likely to be located in promoters and other regulatory regions of genes, rather than in the coding regions. Polymorphisms in the regulatory regions may be particularly relevant in chronic metabolic diseases, such as FCHL, leading to a gradual accumulation of damage over many years before reaching a critical threshold. It is thus possible that, in complex diseases, variations in regulatory sequences cause slight differences in message levels, timing and tissue specificity of gene expression, and protein stability, that lead to subtle quantitative differences in phenotype (Mackay 2001). Accordingly, our finding identifying a novel regulatory element within intron 7 of *USF1* may turn out to exemplify this by possibly providing a novel promoter for a mini-*USF1* protein lacking the transactivation domain (Shoulders 2004).

To conclude, several loci affecting plasma HDL-C and TGs levels have been identified in this thesis (*APOA2* and *APOA1C3A4* gene cluster loci, 8q23, 16q24.1-24.2, and 20q13.11 and 10q11) (I-III). Most interestingly, one of the studies led to the characterization of a new susceptibility gene for FCHL, *USF1* (IV). The strong association between the *USF1* gene and FCHL suggests that *USF1* contributes to the susceptibility to FCHL. Because the *USF1* gene encodes a transcription factor known to regulate several genes participating in glucose and lipid metabolism (for refs. see below), it is potentially the missing link between FCHL and type 2 DM. *USF1* may form a prime candidate in the 1q21-23 region for increasing the susceptibility to FCHL, type 2 DM, and the metabolic syndrome. Moreover, *USF1* might play an important role in low HDL-C, because it resides on chromosome 1 in the vicinity of *APOA2* and the low HDL-C trait was linked to that region (I). Although additional studies are needed to clarify the mechanisms of the connection between *USF1* alleles and the FCHL phenotype, our results provide novel insights into the genetic background of FCHL and low HDL-C, the most common familial lipid disorders predisposing to CHD.

USF1

USF1 is a transcription factor of the basic helix-loop-helix leucine zipper family. It forms homo- and heterodimers (with USF2) and recognizes a CACGTG motif termed E box (Casado et al. 1999; Ribeiro et al. 1999), resulting in activation of gene transcription (Chen et al. 2001) and enhanced expression in response to various stimuli, such as glucose and dietary carbohydrates (Nowak et al. 2004). USF1 is known to regulate several genes participating in glucose and lipid metabolism. In the liver, USF1 regulates the expression of several genes, including fatty acid synthase (FAS) (Casado et al. 1999), APOA2 (Ribeiro et al. 1999) APOC3 (Pastier et al. 2002), APOE (Salero et al. 2003), HL (Botma et al. 2001), and ABCA1 (Yang et al. 2002). In addition, the gene expression of an important factor in hepatic glucose sensing, glucokinase, is regulated by USF1 (Iynedjian 1998). In the pancreas, USF1 controls the expression of insulin (Read et al. 1993), the glucagon receptor (Portois et al. 2002), and the islet-specific glucose-6-phosphatase catalytic-subunit-related gene (Martin et al. 2003a). In adipose tissue, USF1 mediates the insulin-responsive expression of FAS (Wang and Sul 1995; Wang and Sul 1997), the glucose-regulated expression of hormone sensitive lipase (Smih et al. 2002), as well as the gene expression of acetyl CoA carboxylase (Travers et al. 2001). On the basis of these studies, USF1 seems to play a central role in the regulation of several of the key genes of lipid and glucose metabolism.

In the USF knock out mice, the normal transcriptional response of liver genes to glucose seemed to require the presence of either USF1/USF2 heterodimers or USF2 homodimers (Vallet et al. 1998). In these mice, induction of the FAS gene was severely delayed by refeeding a carbohydrate-rich diet, suggesting that USF transactivators, especially USF1/USF2 heterodimers, are essential for sustaining the dietary induction of the FAS gene in the liver (Casado et al. 1999). USF1/USF2 is also capable of activating the transcription of the gene encoding renin, which is an important regulator of the systemic blood pressure (Pan et al. 2001). It has also been proposed that USF genes may play an important role in the differentiation capability of vascular smooth muscle cells, because USF factors modulate the expression of genes that are acutely regulated by the differentiation status of the smooth muscle cells of the vasculature (Chen et al. 2001).

In our study (IV), a specific haplotype of USF1 was associated with FCHL. Interestingly, Putt and colleagues studied three intronic USF1 SNPs in more than 800 healthy men, and reported that the 475C/T-1738C/T haplotype influences the glucose levels during the oral glucose tolerance test (Putt et al. 2004). Furthermore, they found case:control heterogeneity in the interaction between USF1 SNPs genotypes and BMI on fasting LDL and glucose.

It has been suggested that the novel regulatory element identified within intron 7 of USF1 may provide a novel promoter for a mini-USF1 protein lacking the transactivation domain (Shoulders 2004). The putative promoter identified (IV) might operate in vivo to downregulate USF1 activity, because mini-USF proteins have been shown to act in vitro as transdominant inhibitors (Lefrancois-Martinez et al. 1995; Viollet et al. 1996). This theory proposes that intronic variants affecting splice sites or transcription factor binding sites may be of significant importance in complex traits, and this very interesting possibility remains to be investigated.

CONCLUDING REMARKS

In the course of this study, methods for mapping and identifying genes for complex traits have developed tremendously. Especially, the sequence of the human genome produced by the HGP and the genetic databases publicly available for everyone in the World Wide Web are valuable tools for researchers, and are rapidly changing the course of gene hunting from the old methods to a new direction. The increasing numbers of SNPs will facilitate the construction of haplotypes in different chromosomal regions and different populations. Knowledge of the haplotype structures will help to evaluate the association between a genetic marker and a trait in LD mapping. It might also help us to solve which type of populations would be ideal for determining the genetic background of complex disorders.

Having the anatomy of the human genome in front of us offers magnificent opportunities for dissection of the molecular basis of human diseases. Now, in the genomic era, biomedical research is facing changes in its methods and strategies. Structural genomics will focus on functional genomics and proteomics. As we turn from monogenic disorders more and more to dissection of multifactorial disorders, we are monitoring susceptibility rather than causality where analysis of one gene is changing to analysis of multiple genes or gene families, pathways, and systems. Other changes include shifting from research on gene action to gene regulation and from etiology (specific mutation) to pathogenesis (the mechanism of the disease).

Although we have now gained a huge amount of new information about genomics, we do not yet have good guidelines or experience on how to use this knowledge effectively. To get the most out of this new information, efficient collaboration between research groups, data sharing, and pooled data analyses are required in the genomic era. In addition, collaborations among clinicians, epidemiologists, geneticists, statisticians and other experts have become increasingly important and are needed to solve the complex genetic background of multifactorial traits.


Although complex traits are clinically important and common in the population, relatively little is known about the nature of the genetic variation underlying complex diseases. Moreover, most of our knowledge about the genetics of common diseases relates to rare families segregating high-risk alleles (Botstein and Risch 2003), which are generally very infrequent in the population and therefore explain relatively little of overall disease prevalence. In contrast, an important contribution of common variants to the susceptibility to common diseases has been suggested: there are probably many common variants in the human genome with modest but real effects on complex disease risk (Figure 11) (Lohmueller et al. 2003). Likewise, the distribution of quantitative traits in the population is presumed to reflect the cumulative contribution of multiple common DNA sequence variants, each having a small effect (Figure 11).

For complex disorders many chromosomal loci have been identified, but only a small proportion of these findings have led to actual identification of the underlying gene. How can we improve gene identification strategies for complex traits? The tools for the genetic dissection of such traits are improving, and the new methods may even help to better reveal the minor effects of multiple genes in complex disorders. For instance, microarray

methodology provides valuable information about new metabolic pathways, helps to find the genes that are important in disease pathogenesis, and may even show which genes and alleles cause significant risks in the general population. Genome wide association studies, dense SNP maps, and haploblocks of the HapMap-project are examples of tools that most likely will facilitate identification of the genetic background of complex traits.

Figure 11.

A schematic overview of how disease susceptibility alleles, which are common alleles in the population, predispose to a disease. Risk alleles of many different susceptibility genes together with environmental factors cause the disease.



The HGP have made a tremendous contribution to the positional cloning of complex traits by providing a complete catalogue of all the genes residing in a relevant region. Now researchers can go to a variety of websites (such as the National Center for Biotechnology Information, NCBI, <http://www.ncbi.nih.gov> or UCSC Human Genome Browser <http://www.genome.ucsc.edu>) to find tens of thousands of well-characterized genes and to start systematically evaluating candidate genes for the linked chromosomal region instead of tedious original positional cloning.

The most attractive finding of this thesis is the identification of, a susceptibility gene for FCHL in Finnish families, USF1. Nevertheless, this study also exemplifies many problems encountered in the genetic research of complex traits. Probably the most typical question is how to prove that noncoding variants that are associated with a trait really predispose to disease? It seems that not only one SNP but rather a group of adjacent SNPs residing in a relatively wide chromosomal region are associated with disease due to strong LD between these SNPs. So which one is *the* variant and *the* gene that causes susceptibility to disease?

To understand the mechanisms of USF1 in FCHL and to clarify whether USF1 also plays a role in the metabolic syndrome, additional studies are needed. The future plans should include several different approaches. First of all, the challenge ahead is to confirm our promising genetic data through replication. Investigation of the associated USF1 variants in the Dutch FCHL families and in large epidemiological population samples such as the Finnish cohort “Finrisk 1992” is currently ongoing. A recent study by Putt and colleagues offered further evidence that USF1 influences glucose and lipid homeostasis, even in healthy men (Putt et al. 2004).

Second, functional studies of the associated USF1 SNPs, especially *usf1s1* and *usf1s2*, are warranted to determine the roles of the different isoforms of USF1 in the regulation of genes important for lipid and glucose homeostasis. For example, functional proof of regulatory sequence variations may be achieved by techniques of cell biology or bacterial artificial chromosome transgenesis (Symula et al. 1999).

Third, further microarray studies in a larger study sample are needed in order to be able to confirm our preliminary results. The microarray studies should be carried out not only in fat tissue samples but also in different tissues, for instance in muscle tissue obtained from muscle biopsies.

Fourth, animal studies could help us to better understand the underlying molecular mechanism of the USF1 gene in FCHL. For example a transgenic USF1 mouse model could offer a useful tool for that.

Fifth, in order to analyze the role of the USF1 gene in type 2 DM, investigations of the associated USF1 variants in type 2 DM study samples are necessary. In summary, further studies in large epidemiological samples of the Finnish population, and in other well-defined FCHL samples from more mixed populations, as well as in type 2 DM families, will help to clarify the contribution of USF1 not only to FCHL but also to other FCHL-associated phenotypes, type 2 DM and the metabolic syndrome.

FCHL, the most common inherited disorder of abnormal blood lipid levels, is a highly atherogenic multifactorial disorder. As in other complex traits, multiple susceptibility and modifying genes are believed to play a role in the pathogenesis of FCHL. The identified USF1 gene forms a promising candidate gene for FCHL and is also an encouraging example of a gene detected for a complex multifactorial lipid trait. However, the major task in future projects is not only to confirm the role of USF1 but also to identify the other genes associated with FCHL.

In Finnish low HDL-C families, gene hunting will continue. The impact of the USF1 gene in low HDL-C families needs to be investigated. In addition, the role of the APOA2 gene and APOA1C3A4A5 gene cluster in low HDL-C requires additional studies. Future plans include further investigation of the loci linked to the low HDL-C trait found in our previous study (8q23, 16q24.1-24.2 and 20q13.11), as well as sequencing of the promising candidate genes located on 10q11, a region associated with HDL-C and TG levels. The idea of the haplotype blocks capturing most of the genetic variation in a region could be employed as a new strategy to identify the gene for HDL-C on 10q11. If the 10q

region fits the block theory well, this novel approach could significantly decrease the number of SNPs that need to be analysed in order to find the associated gene.

The genomic era offers us great opportunities and only the future will show us, how well we will be able to take advantage of the huge amount of information now available about the human genome. In the end, the ultimate goal is to translate genomic research into health applications, i.e. to harness genetics to help find better ways of diagnosing, treating and preventing disease, and finally, to improve health.

ACKNOWLEDGEMENTS

This study was carried out in the Department of Molecular Medicine at the National Public Health Institute, Helsinki, Finland and UCLA, Department of Human Genetics, Gonda Neuroscience and Genetics Research Center, Los Angeles, USA during 1998-2004. I wish to thank Jussi Huttunen and Pekka Puska, the former and present heads of the National Public Health institute for providing excellent research facilities.

This study was financially supported by grants from the University of Helsinki, the Academy of Finland (Center of Excellence on Disease Genetics), the Ulla Hjelt Fund of the Foundation for Pediatric Research, Aarne Koskelo Foundation, Maud Kuistila Foundation, Paulo Foundation, Finnish Cultural Foundation, Biomedicum Helsinki Foundation, Helsinki University Central Hospital Research Foundation, Finnish Foundation for Cardiovascular Research, as well as the Finnish Medical Foundation. They are all gratefully acknowledged.

I am grateful to the Helsinki Biomedical Graduate School (HBGS) for giving me the opportunity to begin the MD/PhD program already during the first year of my medical studies. I also wish to thank HBGS for providing financial support in addition to the scientific training and tuition.

I want to express my deepest gratitude to my excellent supervisors, Professor Leena Palotie and Assistant Professor Päivi Pajukanta. I'm impressed by Leena's devotion to science, her endless energy and effectiveness, as well as her inspiring spirit. I am especially grateful for Leena not only for the good tutoring and great research opportunities of her laboratory, but also for providing me the possibility to do my PhD project side by side with my medical studies. I also thank her for the great opportunity to work for a year in UCLA.

I owe a large debt of gratitude to my other supervisor, Päivi Pajukanta. Päivi has held a key position in this project, and her tremendous knowledge in genetics on complex lipid traits has been extremely valuable and even irreplaceable during the course of this study, she has taught me a lot. Päivi's enthusiasm towards science is inspiring. Päivi, I sincerely thank you for your patience and guidance throughout all of these years. Thanks for sharing your knowledge and skills with me and for being the driving force behind this thesis.

My sincere thanks go to our collaborator, Professor Marja-Riitta Taskinen, Marsu. She has greatly contributed to this study, not only by providing the unique dyslipidemic family material for our genetic studies, but also by her enormous knowledge on lipid disorders and diabetes. I greatly respect her successful career as a doctor, and especially the way she has been able to combine clinical and scientific work.

Professor Kimmo Kontula and Docent Matti Jauhiainen, the senior members of my thesis committee, are warmly thanked for their positive feedback and constructive comments. I greatly admire Matti's enthusiasm for science and his excellent knowledge on lipoprotein metabolism. Professor Petri Kovanen and Docent Katriina Aalto-Setälä are also warmly thanked for reviewing the thesis manuscript and for their valuable comments and criticisms.

I am grateful to Professor Christian Ehnholm and to the senior scientists of the lab, Anu Jalanko and Ismo Ulmanen, Iski. Thank you for maintaining the lab at a high standard while Leena was in UCLA. Christian is acknowledged for his interest in and for his contribution towards HDL-C metabolism research. Anu and Iski are thanked for including me into many of the social events

within their own research teams. Iski is the father figure of the lab, the one who always asks you if everything is OK when you meet him in the corridor, and he is a person with whom you can talk to if something weighs upon you. Iski is also thanked for many special events he has arranged outside the lab. I have really enjoyed the nature trips, bird watching and mushroom picking, to name but a few.

My sincere thanks go to Aino Soro-Paavonen for her friendship and active collaboration. She has provided pleasant company during our 'business travels', accompanied me jogging during the early hours of the mornings, and offered great pig-parties in Loimaa as well as many other unforgettable moments together. Most of all, I want to thank you Aino for the enormous work you have done with the HDL- and FCHL-family collection, the phenotypes, and the genome-wide scan. Kati Ylitalo is also acknowledged for her contribution to the family collection.

I want to thank my closest co-worker Elina Suviolahti for making the last years in lab more enjoyable. I was fortunate to have you in the same project. Once you arrived I wasn't alone anymore as we could share all of the problems and adversities of the research project together. Elina, I love your eastern-Finnish style of speaking, your positive attitude and your endless enthusiasm for surfing and skiing.

Markus Perola is thanked for his mentorship, his support, and for his nice jokes every now and then. Markus has guided and helped me in many ways when both of my supervisors were in LA. I also want to extend my thanks to Tero Hiekkalinna, who has helped and assisted me many times with data analyses and computer problems. Many thanks also to Pekka Ellonen for his help, good tips, and technical support. I am happy that we have such a skillful and talented guy like you Pekka in our lab.

Minna Levander, Päivi Tainola, Arja Terola, Arja Tapio, and Elli Kempas are acknowledged for their technical assistance. Elli is especially thanked for her help and guidance in the lab when I first started. Thanks also to the ex-DNA-master, Mervi Alanne, and to our new one Outi Repola.

I am immensely grateful to all of my workmates in the lab. Thanks for making those long days in lab easier to bare. A big thanks to you all, especially: Jussi Naukkarinen, Jenny Ekholm, Mira Kyttälä, Henna Haravuori, Sampo Sammalisto, Maria Halonen, Laura Ahtiainen, and Tanja Ilmarinen. I also wish to thank Anna, Tintti, Tony, Denis, Kismat, Emma, Nora, Taina, Annina, Janna, Will, Ilona, Mari, Tero, Mikko, among many others, for their great company.

Ville Holmberg is thanked for his friendship in the office and for all of those interesting conversations and active 'global' discussions. Juha Paloneva and Niklas Pakkasjärvi were great company and offered many cheerful moments. Many thanks go to Jenni Leppävuori for her positive attitude, for all those sporty and funny moments, and for sharing Leena's house with us. I wish to thank Aimee Trudeau for sharing the office with me and for always being ready to help me with any language problems, as well as to Sarah Siggins who is also thanked for her good company especially in the conference trips because without you the meeting in Kyoto would not have been that cheerful.

I had a great time in UCLA and I want to thank everyone there, especially Outi Saranen, Susanna Vikman, Jenny, Chris, Greg, Tam, Ed, Leena, Nikke&Leila, Juha&Petra, Mira, Hilde, Jenni and Aino. Most of all I'd like to thank Kisse for sharing the year in UCLA with me, such as our flat in Lindbrook Manor, for being a part of our small 'family', for the trips to Yosemite, San Fransisco, Mexico, for everyday life with its ups and downs and for moments with Tempukka. I still miss you at times and those cheerful moments during our stay in LA.

My sincere thanks go to Kaisu Luiro and Joni Turunen. Thanks for the friendship and shared moments in the lab, med school and beyond. I love Joni's enthusiastic attitude to science, dancing, partying, shopping - whatever it happens to be. Kaisu, thanks for the tea-breaks, for Yahoo-messaging, sharing the ups and downs that come with Ph.D. work, for the laughs and for the conversations about difficult things in life. I wish to thank also other friends from Cursus Metamorphosis; Hanna & Teemu, Jenni, Sampo, Kipa, Sole, Jaakko, Tiina, Tarmo, Nuutti, Birgi, and Jonna: thanks for the great company during med school and thereafter, the enjoyable summer cottage trips, crab parties and many funny moments. Laura Kotaniemi thanks for your friendship, fascinating talks about everything and good, wise opinions. Thanks also to the students of Cursus Unicus for being the second cursus of mine after I came back from UCLA.

Many thanks to my high-school friends from Mäkelänrinne, especially Karo, Eira, Miia, Taru and Mirva. Special thanks go to my 'personal shrink' Sevi for being there in times of desperation. Thanks also to my old friend Mia - after all, we found our way to sunshine and I think we did well! Many beers (with lemon and ice) are owed to my cousin Sanna for always being ready to party with me.

My dear friend Kati Koskela is thanked for her positive energy and for joyful times together. Our long runs and walks to Seurasaari have been essential for me after long days in the lab and endless dates with the pipette or computer. We have shared so many moments of happiness and tears, ups and downs of life. Thanks for the countless crazy laughs but most of all, thanks for offering a shoulder to cry on when most needed.

Michel, I'd like to thank you for the love and support you gave me and for sharing your life with me during many years in medical school and during this thesis. I miss u and the good times together. In pain we grow.

I am grateful to my grandparents, Eino and Eevi, and my late grandmother Tuija and grandpa Viljo. Tuija was the heart of our clan, and I thank my grandfather Eino for his support and for constantly showing interest in my studies. My grandmother Eevi is also thanked for always being so proud of me.

I owe my warmest thanks to my parents Maisa and Eero for their care and support throughout my life. I am grateful that their love has never been dependent on my achievements. Arja is not only thanked for all her delicious cooking but also for taking care of my father and my dear little sisters Anri and Anja, and for the love we shared towards Lari, the one and only dog in my life. I wish to thank my twin-brother Jouni for sharing at least the first 15 years of our life with me. It's cool to be a twin! However, sometimes I really wonder about our dissimilarity and I envy your easy-going attitude towards life. Thanks Jouni for bringing such beautiful babies into our family, especially sweet Emma who has become the sunshine of my life. My deepest thanks go to my dear big sister Vippe and her family including Ilkka, Oskari and Veera. With you I have had many happy and relaxing moments, especially in Viinimäki.

Finally, I want to express my sincere thanks to the patients who have participated in this study. Without their involvement this study would not have been possible.

Helsinki, December 2004

Heidi Lilja

REFERENCES

- Acton S, Osgood D, Donoghue M, Corella D, Pocovi M, Cenarro A, Mozas P, Keilty J, Squazzo S, Woolf EA, Ordovas JM (1999) Association of polymorphisms at the SR-BI gene locus with plasma lipid levels and body mass index in a white population. *Arterioscler Thromb Vasc Biol* 19:1734-43
- Aitman TJ, Godsfand IF, Farren B, Crook D, Wong HJ, Scott J (1997) Defects of insulin action on fatty acid and carbohydrate metabolism in familial combined hyperlipidemia. *Arterioscler Thromb Vasc Biol* 17:748-54
- Alberti KG, Zimmet PZ (1998) Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabet Med* 15:539-53
- Allayee H, Aouizerat BE, Cantor RM, Dallinga-Thie GM, Krauss RM, Lanning CD, Rotter JI, Lusis AJ, de Bruin TW (1998) Families with familial combined hyperlipidemia and families enriched for coronary artery disease share genetic determinants for the atherogenic lipoprotein phenotype. *Am J Hum Genet* 63:577-85
- Allayee H, Castellani LW, Cantor RM, de Bruin TW, Lusis AJ (2003) Biochemical and genetic association of plasma apolipoprotein A-II levels with familial combined hyperlipidemia. *Circ Res* 92:1262-7
- Allayee H, Dominguez KM, Aouizerat BE, Krauss RM, Rotter JI, Lu J, Cantor RM, de Bruin TW, Lusis AJ (2000) Contribution of the hepatic lipase gene to the atherogenic lipoprotein phenotype in familial combined hyperlipidemia. *J Lipid Res* 41:245-52
- Allayee H, Krass KL, Pajukanta P, Cantor RM, van der Kallen CJ, Mar R, Rotter JI, de Bruin TW, Peltonen L, Lusis AJ (2002) Locus for elevated apolipoprotein B levels on chromosome 1p31 in families with familial combined hyperlipidemia. *Circ Res* 90:926-31
- Almasy L, Blangero J (1998) Multipoint quantitative-trait linkage analysis in general pedigrees. *Am J Hum Genet* 62:1198-211
- Almasy L, Hixson JE, Rainwater DL, Cole S, Williams JT, Mahaney MC, VandeBerg JL, Stern MP, MacCluer JW, Blangero J (1999) Human pedigree-based quantitative-trait-locus mapping: localization of two genes influencing HDL-cholesterol metabolism. *Am J Hum Genet* 64:1686-93
- Amos AF, McCarty DJ, Zimmet P (1997) The rising global burden of diabetes and its complications: estimates and projections to the year 2010. *Diabet Med* 14 Suppl 5:S1-85
- Amos CI (1994) Robust variance-components approach for assessing genetic linkage in pedigrees. *Am J Hum Genet* 54:535-43
- Amos CI, de Andrade M (2001) Genetic linkage methods for quantitative traits. *Stat Methods Med Res* 10:3-25
- Angius A, Melis PM, Morelli L, Petretto E, Casu G, Maestrale GB, Fraumene C, Bebbere D, Forabosco P, Pirastu M (2001) Archival, demographic and genetic studies define a Sardinian sub-isolate as a suitable model for mapping complex traits. *Hum Genet* 109:198-209
- Anzalone DA, Anzalone FL, Fos PJ (1995) High-density lipoprotein-cholesterol: determining hygienic factors for intervention. *J Occup Environ Med* 37:856-61
- Aouizerat BE, Allayee H, Bodnar J, Krass KL, Peltonen L, de Bruin TW, Rotter JI, Lusis AJ (1999a) Novel genes for familial combined hyperlipidemia. *Curr Opin Lipidol* 10:113-22
- Aouizerat BE, Allayee H, Cantor RM, Dallinga-Thie GM, Lanning CD, de Bruin TW, Lusis AJ, Rotter JI (1999b) Linkage of a candidate gene locus to familial combined hyperlipidemia: lecithin:cholesterol acyltransferase on 16q. *Arterioscler Thromb Vasc Biol* 19:2730-6
- Aouizerat BE, Allayee H, Cantor RM, Davis RC, Lanning CD, Wen PZ, Dallinga-Thie GM, de Bruin TW, Rotter JI, Lusis AJ (1999c) A genome scan for familial combined hyperlipidemia reveals evidence of linkage with a locus on chromosome 11. *Am J Hum Genet* 65:397-412
- Aouizerat BE, Kane JP (2003) Apolipoprotein A-II: active or passive role in familial combined hyperlipidemia. *Circ Res* 92:1179-81
- Aouizerat BE, Kulkarni M, Heilbron D, Drown D, Raskin S, Pullinger CR, Malloy MJ, Kane JP (2003) Genetic analysis of a polymorphism in the human apoA-V gene: effect on plasma lipids. *J Lipid Res* 44:1167-73
- Arya R, Duggirala R, Almasy L, Rainwater DL, Mahaney MC, Cole S, Dyer TD, Williams K, Leach RJ, Hixson JE, MacCluer JW, O'Connell P, Stern MP, Blangero J (2002) Linkage of high-density lipoprotein-cholesterol concentrations to a locus on chromosome 9p in Mexican Americans. *Nat Genet* 30:102-5
- Assmann G, Gotto AM, Jr. (2004) HDL cholesterol and protective factors in atherosclerosis. *Circulation* 109:III8-14
- Assmann G, Nofer JR (2003) Atheroprotective effects of high-density lipoproteins. *Annu Rev Med* 54:321-41
- Assmann G, Schulte H, von Eckardstein A, Huang Y (1996) High-density lipoprotein cholesterol as a predictor of coronary heart disease risk. The PROCAM experience and pathophysiological implications for reverse cholesterol transport. *Atherosclerosis* 124 Suppl:S11-20
- Assmann G, von Eckardstein A, Brewer HB (1995) Familial high density lipoprotein deficiency: Tangier disease. McGraw-Hill, New York
- Assmann G, von Eckardstein A, Funke H (1993) High density lipoproteins, reverse transport of cholesterol, and coronary artery disease. Insights from mutations. *Circulation* 87:III28-34

- Auranen M, Vanhala R, Varilo T, Ayers K, Kempas E, Ylisaukko-Oja T, Sinsheimer JS, Peltonen L, Jarvela I (2002) A genomewide screen for autism-spectrum disorders: evidence for a major susceptibility locus on chromosome 3q25-27. *Am J Hum Genet* 71:777-90
- Austin MA, Brunzell JD, Fitch WL, Krauss RM (1990a) Inheritance of low density lipoprotein subclass patterns in familial combined hyperlipidemia. *Arteriosclerosis* 10:520-30
- Austin MA, King MC, Vranizan KM, Krauss RM (1990b) Atherogenic lipoprotein phenotype. A proposed genetic marker for coronary heart disease risk. *Circulation* 82:495-506
- Ayyobi AF, Brunzell JD (2003) Lipoprotein distribution in the metabolic syndrome, type 2 diabetes mellitus, and familial combined hyperlipidemia. *Am J Cardiol* 92:27J-33J
- Babirak SP, Brown BG, Brunzell JD (1992) Familial combined hyperlipidemia and abnormal lipoprotein lipase. *Arterioscler Thromb* 12:1176-83
- Barter P, Kastelein J, Nunn A, Hobbs R (2003) High density lipoproteins (HDLs) and atherosclerosis; the unanswered questions. *Atherosclerosis* 168:195-211
- Barzilai N, Atzmon G, Schechter C, Schaefer EJ, Cupples AL, Lipton R, Cheng S, Shuldiner AR (2003) Unique lipoprotein phenotype and genotype associated with exceptional longevity. *Jama* 290:2030-40
- Baum L, Tomlinson B, Thomas GN (2003) APOA5-1131T>C polymorphism is associated with triglyceride levels in Chinese men. *Clin Genet* 63:377-9
- Blangero J, Williams JT, Almasy L (2001) Variance component methods for detecting complex trait loci. *Adv Genet* 42:151-81
- Bodnar JS, Chatterjee A, Castellani LW, Ross DA, Ohmen J, Cavalcoli J, Wu C, Dains KM, Catanese J, Chu M, Sheth SS, Charugundla K, Demant P, West DB, de Jong P, Lusis AJ (2002) Positional cloning of the combined hyperlipidemia gene *Hyplip1*. *Nat Genet* 30:110-6
- Bodzioch M, Orso E, Klucken J, Langmann T, Bottcher A, Diederich W, Drobnik W, Barlage S, Buchler C, Porsch-Ozcurumez M, Kaminski WE, Hahmann HW, Oette K, Rothe G, Aslanidis C, Lackner KJ, Schmitz G (1999) The gene encoding ATP-binding cassette transporter 1 is mutated in Tangier disease. *Nat Genet* 22:347-51
- Botma GJ, Verhoeven AJ, Jansen H (2001) Hepatic lipase promoter activity is reduced by the C-480T and G-216A substitutions present in the common LIPC gene variant, and is increased by Upstream Stimulatory Factor. *Atherosclerosis* 154:625-32
- Botstein D, Risch N (2003) Discovering genotypes underlying human phenotypes: past successes for mendelian disease, future approaches for complex disease. *Nat Genet* 33 Suppl:228-37
- Bowden DW, Sale M, Howard TD, Qadri A, Spray BJ, Rothschild CB, Akots G, Rich SS, Freedman BI (1997) Linkage of genetic markers on human chromosomes 20 and 12 to NIDDM in Caucasian sib pairs with a history of diabetic nephropathy. *Diabetes* 46:882-6
- Bredie SJ, Tack CJ, Smits P, Stalenhoef AF (1997a) Nonobese patients with familial combined hyperlipidemia are insulin resistant compared with their nonaffected relatives. *Arterioscler Thromb Vasc Biol* 17:1465-71
- Bredie SJ, van Drongelen J, Kiemeny LA, Demacker PN, Beaty TH, Stalenhoef AF (1997b) Segregation analysis of plasma apolipoprotein B levels in familial combined hyperlipidemia. *Arterioscler Thromb Vasc Biol* 17:834-40
- Bredie SJ, Vogelaar JM, Demacker PN, Stalenhoef AF (1996) Apolipoprotein E polymorphism influences lipid phenotypic expression, but not the low density lipoprotein subfraction distribution in familial combined hyperlipidemia. *Atherosclerosis* 126:313-24
- Brewer HB, Jr., Santamarina-Fojo S (2003) Clinical significance of high-density lipoproteins and the development of atherosclerosis: focus on the role of the adenosine triphosphate-binding cassette protein A1 transporter. *Am J Cardiol* 92:10K-16K
- Broeckel U, Hengstenberg C, Mayer B, Holmer S, Martin LJ, Comuzzie AG, Blangero J, Nurnberg P, Reis A, Riegger GA, Jacob HJ, Schunkert H (2002) A comprehensive linkage analysis for myocardial infarction and its related risk factors. *Nat Genet* 30:210-4
- Brooks-Wilson A, Marcel M, Clee SM, Zhang LH, Roomp K, van Dam M, Yu L, et al. (1999) Mutations in ABC1 in Tangier disease and familial high-density lipoprotein deficiency. *Nat Genet* 22:336-45
- Brousseau T, Dupuy-Gorce AM, Evans A, Arveiler D, Ruidavets JB, Haas B, Cambou JP, Luc G, Ducimetiere P, Amouyel P, Helbecque N (2002) Significant impact of the highly informative (CA)_n repeat polymorphism of the APOA-II gene on the plasma APOA-II concentrations and HDL subfractions: The ECTIM study. *Am J Med Genet* 110:19-24
- Brown ML, Inazu A, Hesler CB, Agellon LB, Mann C, Whitlock ME, Marcel YL, Milne RW, Koizumi J, Mabuchi H, et al. (1989) Molecular basis of lipid transfer protein deficiency in a family with increased high-density lipoproteins. *Nature* 342:448-51
- Brown MS, Basu SK, Falck JR, Ho YK, Goldstein JL (1980) The scavenger cell pathway for lipoprotein degradation: specificity of the binding site that mediates the uptake of negatively-charged LDL by macrophages. *J Supramol Struct* 13:67-81
- Brunzell JD, Albers JJ, Chait A, Grundy SM, Groszek E, McDonald GB (1983) Plasma lipoproteins in familial combined hyperlipidemia and monogenic familial hypertriglyceridemia. *J Lipid Res* 24:147-55
- Bu X, Warden CH, Xia YR, De Meester C, Puppione DL, Teruya S, Lokensgard B, Daneshmand S, Brown J, Gray RJ, et al. (1994) Linkage analysis of the genetic determinants of high density lipoprotein concentrations and

- composition: evidence for involvement of the apolipoprotein A-II and cholesteryl ester transfer protein loci. *Hum Genet* 93:639-48
- Bucher KD, Friedlander Y, Kaplan EB, Namboodiri KK, Kark JD, Eisenberg S, Stein Y, Rifkind BM (1988) Biological and cultural sources of familial resemblance in plasma lipids: a comparison between North America and Israel--the Lipid Research Clinics Program. *Genet Epidemiol* 5:17-33
- Campagna F, Montali A, Baroni MG, Maria AT, Ricci G, Antonini R, Verna R, Arca M (2002) Common variants in the lipoprotein lipase gene, but not those in the insulin receptor substrate-1, the beta3-adrenergic receptor, and the intestinal fatty acid binding protein-2 genes, influence the lipid phenotypic expression in familial combined hyperlipidemia. *Metabolism* 51:1298-305
- Carlson CS, Eberle MA, Kruglyak L, Nickerson DA (2004) Mapping complex disease loci in whole-genome association studies. *Nature* 429:446-52
- Carr MC, Brunzell JD (2004) Abdominal obesity and dyslipidemia in the metabolic syndrome: importance of type 2 diabetes and familial combined hyperlipidemia in coronary artery disease risk. *J Clin Endocrinol Metab* 89:2601-7
- Casado M, Vallet VS, Kahn A, Vaulont S (1999) Essential role in vivo of upstream stimulatory factors for a normal dietary response of the fatty acid synthase gene in the liver. *J Biol Chem* 274:2009-13
- Castellani LW, Weinreb A, Bodnar J, Goto AM, Doolittle M, Mehrabian M, Demant P, Lusis AJ (1998) Mapping a gene for combined hyperlipidaemia in a mutant mouse strain. *Nat Genet* 18:374-7
- Castelli WP, Doyle JT, Gordon T, Hames CG, Hjortland MC, Hulley SB, Kagan A, Zukel WJ (1977) HDL cholesterol and other lipids in coronary heart disease. The cooperative lipoprotein phenotyping study. *Circulation* 55:767-72
- Castelli WP, Garrison RJ, Wilson PW, Abbott RD, Kalousdian S, Kannel WB (1986) Incidence of coronary heart disease and lipoprotein cholesterol levels. The Framingham Study. *Jama* 256:2835-8
- Castro Cabezas M, de Bruin TW, de Valk HW, Shoulders CC, Jansen H, Willem Erkelens D (1993) Impaired fatty acid metabolism in familial combined hyperlipidemia. A mechanism associating hepatic apolipoprotein B overproduction and insulin resistance. *J Clin Invest* 92:160-8
- Cederberg A, Gronning LM, Ahren B, Tasken K, Carlsson P, Enerback S (2001) FOXC2 is a winged helix gene that counteracts obesity, hypertriglyceridemia, and diet-induced insulin resistance. *Cell* 106:563-73
- Chakravarthy MV, Booth FW (2004) Eating, exercise, and "thrifty" genotypes: connecting the dots toward an evolutionary understanding of modern chronic diseases. *J Appl Physiol* 96:3-10
- Chakravarti A (2001) To a future of genetic medicine. *Nature* 409:822-3
- Chen YH, Layne MD, Watanabe M, Yet SF, Perrella MA (2001) Upstream stimulatory factors regulate aortic preferentially expressed gene-1 expression in vascular smooth muscle cells. *J Biol Chem* 276:47658-63
- Cheng JF, Pennacchio LA (2003) Comparative and functional analysis of cardiovascular-related genes. *Pharmacogenomics* 4:571-82
- Chinetti G, Gbaguidi FG, Griglio S, Mallat Z, Antonucci M, Poulain P, Chapman J, Fruchart JC, Tedgui A, Najib-Fruchart J, Staels B (2000) CLA-1/SR-BI is expressed in atherosclerotic lesion macrophages and regulated by activators of peroxisome proliferator-activated receptors. *Circulation* 101:2411-7
- Chittur SV (2004) DNA microarrays: tools for the 21st Century. *Comb Chem High Throughput Screen* 7:531-7
- Claverie JM (2001) Gene number. What if there are only 30,000 human genes? *Science* 291:1255-7
- Clerget-Darpoux F, Bonaiti-Pellie C, Hochez J (1986) Effects of misspecifying genetic parameters in lod score analysis. *Biometrics* 42:393-9
- Cohen JC, Kiss RS, Pertsemlidis A, Marcel YL, McPherson R, Hobbs HH (2004) Multiple rare alleles contribute to low plasma levels of HDL cholesterol. *Science* 305:869-72
- Cohen JC, Wang Z, Grundy SM, Stoesz MR, Guerra R (1994) Variation at the hepatic lipase and apolipoprotein AI/III/AIV loci is a major cause of genetically determined variation in plasma HDL cholesterol levels. *J Clin Invest* 94:2377-84
- Collins FS (1995) Positional cloning moves from perdictional to traditional. *Nat Genet* 9:347-50
- Collod-Beroud G, Boileau C (2002) Marfan syndrome in the third Millennium. *Eur J Hum Genet* 10:673-81
- Coon H, Myers RH, Borecki IB, Arnett DK, Hunt SC, Province MA, Djousse L, Leppert MF (2000) Replication of linkage of familial combined hyperlipidemia to chromosome 1q with additional heterogeneous effect of apolipoprotein A-I/C-III/A-IV locus. The NHLBI Family Heart Study. *Arterioscler Thromb Vasc Biol* 20:2275-80
- Coon H, Singh N, Dunn D, Eckfeldt JH, Province MA, Hopkins PN, Weiss R, Hunt SC, Leppert MF (2004) TXNIP gene not associated with familial combined hyperlipidemia in the NHLBI Family Heart Study. *Atherosclerosis* 174:357-62
- Criqui MH, Wallace RB, Heiss G, Mishkel M, Schonfeld G, Jones GT (1980) Cigarette smoking and plasma high-density lipoprotein cholesterol. The Lipid Research Clinics Program Prevalence Study. *Circulation* 62:IV70-6
- Cross HE (1976) Population studies and the Old Order Amish. *Nature* 262:17-20
- Curtis D, Sham PC, Vallada HP (1995) Genetic analysis of complex disease. *Nat Genet* 9:13
- Cuthbert JA, East CA, Bilheimer DW, Lipsky PE (1986) Detection of familial hypercholesterolemia by assaying functional low-density-lipoprotein receptors on lymphocytes. *N Engl J Med* 314:879-83

- Dahlman I, Eaves IA, Kosoy R, Morrison VA, Heward J, Gough SC, Allahabadia A, Franklyn JA, Tuomilehto J, Tuomilehto-Wolf E, Cucca F, Guja C, Ionescu-Tirgoviste C, Stevens H, Carr P, Nutland S, McKinney P, Shield JP, Wang W, Cordell HJ, Walker N, Todd JA, Concannon P (2002) Parameters for reliable results in genetic association studies in common disease. *Nat Genet* 30:149-50
- Dallinga-Thie GM, Bu XD, van Linde-Sibenius Trip M, Rotter JI, Lusis AJ, de Bruin TW (1996) Apolipoprotein A-I/C-III/A-IV gene cluster in familial combined hyperlipidemia: effects on LDL-cholesterol and apolipoproteins B and C-III. *J Lipid Res* 37:136-47
- Dallinga-Thie GM, van Linde-Sibenius Trip M, Rotter JI, Cantor RM, Bu X, Lusis AJ, de Bruin TW (1997) Complex genetic contribution of the Apo AI-CIII-AIV gene cluster to familial combined hyperlipidemia. Identification of different susceptibility haplotypes. *J Clin Invest* 99:953-61
- Daly MJ, Rioux JD, Schaffner SF, Hudson TJ, Lander ES (2001) High-resolution haplotype structure in the human genome. *Nat Genet* 29:229-32
- Damcott CM, Sack P, Shuldiner AR (2003) The genetics of obesity. *Endocrinol Metab Clin North Am* 32:761-86
- de Bruin TW, Mailly F, van Barlingen HH, Fisher R, Castro Cabezas M, Talmud P, Dallinga-Thie GM, Humphries SE (1996) Lipoprotein lipase gene mutations D9N and N291S in four pedigrees with familial combined hyperlipidaemia. *Eur J Clin Invest* 26:631-9
- de Graaf J, Stalenhoef AF (1998) Defects of lipoprotein metabolism in familial combined hyperlipidaemia. *Curr Opin Lipidol* 9:189-96
- de la Chapelle A (1993) Disease gene mapping in isolated human populations: the example of Finland. *J Med Genet* 30:857-65
- Deeb SS, Nevin DN, Iwasaki L, Brunzell JD (1996) Two novel apolipoprotein A-IV variants in individuals with familial combined hyperlipidemia and diminished levels of lipoprotein lipase activity. *Hum Mutat* 8:319-25
- Deeb SS, Takata K, Peng RL, Kajiyama G, Albers JJ (1990) A splice-junction mutation responsible for familial apolipoprotein A-II deficiency. *Am J Hum Genet* 46:822-7
- Deeb SS, Zambon A, Carr MC, Ayyobi AF, Brunzell JD (2003) Hepatic lipase and dyslipidemia: interactions among genetic variants, obesity, gender, and diet. *J Lipid Res* 44:1279-86
- DeFronzo RA (1997) Insulin resistance: a multifaceted syndrome responsible for NIDDM, obesity, hypertension, dyslipidaemia and atherosclerosis. *Neth J Med* 50:191-7
- Demant T, Carlson LA, Holmquist L, Karpe F, Nilsson-Ehle P, Packard CJ, Shepherd J (1988) Lipoprotein metabolism in hepatic lipase deficiency: studies on the turnover of apolipoprotein B and on the effect of hepatic lipase on high density lipoprotein. *J Lipid Res* 29:1603-11
- Devroey D, De Swaef N, Coigniez P, Vandevoorde J, Kartounian J, Betz W (2004) Correlations between lipid levels and age, gender, glycemia, obesity, diabetes, and smoking. *Endocr Res* 30:83-93
- Durstine JL, Grandjean PW, Davis PG, Ferguson MA, Alderson NL, DuBose KD (2001) Blood lipid and lipoprotein adaptations to exercise: a quantitative analysis. *Sports Med* 31:1033-62
- Eichenbaum-Voline S, Olivier M, Jones EL, Naoumova RP, Jones B, Gau B, Patel HN, Seed M, Betteridge DJ, Galton DJ, Rubin EM, Scott J, Shoulders CC, Pennacchio LA (2004) Linkage and association between distinct variants of the APOA1/C3/A4/A5 gene cluster and familial combined hyperlipidemia. *Arterioscler Thromb Vasc Biol* 24:167-74
- Eisenberg S (1984) High density lipoprotein metabolism. *J Lipid Res* 25:1017-58
- Ekelund J, Lichtermann D, Hovatta I, Ellonen P, Suvisaari J, Terwilliger JD, Juvonen H, Varilo T, Arajärvi R, Kokko-Sahin ML, Lonnqvist J, Peltonen L (2000) Genome-wide scan for schizophrenia in the Finnish population: evidence for a locus on chromosome 7q22. *Hum Mol Genet* 9:1049-57
- Ekholm JM, Kieseppa T, Hiekkalinna T, Partonen T, Paunio T, Perola M, Ekelund J, Lonnqvist J, Pekkarinen-Ijas P, Peltonen L (2003) Evidence of susceptibility loci on 4q32 and 16p12 for bipolar disorder. *Hum Mol Genet* 12:1907-15
- Elbein SC, Hoffman MD, Teng K, Leppert MF, Hasstedt SJ (1999) A genome-wide search for type 2 diabetes susceptibility genes in Utah Caucasians. *Diabetes* 48:1175-82
- Ellison RC, Zhang Y, Qureshi MM, Knox S, Arnett DK, Province MA (2004) Lifestyle determinants of high-density lipoprotein cholesterol: the National Heart, Lung, and Blood Institute Family Heart Study. *Am Heart J* 147:529-35
- Endo K, Yanagi H, Araki J, Hirano C, Yamakawa-Kobayashi K, Tomura S (2002) Association found between the promoter region polymorphism in the apolipoprotein A-V gene and the serum triglyceride level in Japanese schoolchildren. *Hum Genet* 111:570-2
- Eriksson AW (1973) Genetic polymorphisms in Finno-Ugrian populations. Finns, Lapps and Maris. *Isr J Med Sci* 9:1156-70
- Falk CT, Rubinstein P (1987) Haplotype relative risks: an easy reliable way to construct a proper control sample for risk calculations. *Ann Hum Genet* 51 (Pt 3):227-33
- Franceschini G, Sirtori CR, Capurso A, 2nd, Weisgraber KH, Mahley RW (1980) A-Milano apoprotein. Decreased high density lipoprotein cholesterol levels with significant lipoprotein modifications and without clinical atherosclerosis in an Italian family. *J Clin Invest* 66:892-900
- Friedewald WT, Levy RI, Fredrickson DS (1972) Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 18:499-502

- Friedman TB, Liang Y, Weber JL, Hinnant JT, Barber TD, Winata S, Arhya IN, Asher JH, Jr. (1995) A gene for congenital, recessive deafness DFNB3 maps to the pericentromeric region of chromosome 17. *Nat Genet* 9:86-91
- Froguel P, Velho G (2001) Genetic determinants of type 2 diabetes. *Recent Prog Horm Res* 56:91-105
- Funke H, von Eckardstein A, Pritchard PH, Albers JJ, Kastelein JJ, Droste C, Assmann G (1991) A molecular defect causing fish eye disease: an amino acid exchange in lecithin-cholesterol acyltransferase (LCAT) leads to the selective loss of alpha-LCAT activity. *Proc Natl Acad Sci U S A* 88:4855-9
- Gabriel SB, Schaffner SF, Nguyen H, Moore JM, Roy J, Blumenstiel B, Higgins J, DeFelice M, Lochner A, Faggart M, Liu-Cordero SN, Rotimi C, Adeyemo A, Cooper R, Ward R, Lander ES, Daly MJ, Altshuler D (2002) The structure of haplotype blocks in the human genome. *Science* 296:2225-9
- Gagne E, Genest J, Jr., Zhang H, Clarke LA, Hayden MR (1994) Analysis of DNA changes in the LPL gene in patients with familial combined hyperlipidemia. *Arterioscler Thromb* 14:1250-7
- Gagnon F, Jarvik GP, Motulsky AG, Deeb SS, Brunzell JD, Wijsman EM (2003) Evidence of linkage of HDL level variation to APOC3 in two samples with different ascertainment. *Hum Genet* 113:522-33
- Ganong WF (1997) Review of Medical Physiology, 18th ed., Appelton & Lange, Stamford, Connecticut, USA
- Garcia MJ, McNamara PM, Gordon T, Kannel WB (1974) Morbidity and mortality in diabetics in the Framingham population. Sixteen year follow-up study. *Diabetes* 23:105-11
- Garrison RJ, Kannel WB, Feinleib M, Castelli WP, McNamara PM, Padgett SJ (1978) Cigarette smoking and HDL cholesterol: the Framingham offspring study. *Atherosclerosis* 30:17-25
- Gehrisch S, Kostka H, Tiebel M, Patzak A, Paetzold A, Julius U, Schroeder HE, Hanefeld M, Jaross W (1999) Mutations of the human hepatic lipase gene in patients with combined hypertriglyceridemia/hyperalphalipoproteinemia and in patients with familial combined hyperlipidemia. *J Mol Med* 77:728-34
- Genest JJ, Jr., Martin-Munley SS, McNamara JR, Ordovas JM, Jenner J, Myers RH, Silberman SR, Wilson PW, Salem DN, Schaefer EJ (1992) Familial lipoprotein disorders in patients with premature coronary artery disease. *Circulation* 85:2025-33
- Geurts JM, Janssen RG, van Greevenbroek MM, van der Kallen CJ, Cantor RM, Bu X, Aouizerat BE, Allayee H, Rotter JI, de Bruin TW (2000) Identification of TNFRSF1B as a novel modifier gene in familial combined hyperlipidemia. *Hum Mol Genet* 9:2067-74
- Ghosh S, Watanabe RM, Hauser ER, Valle T, Magnuson VL, Erdos MR, Langefeld CD, et al. (1999) Type 2 diabetes: evidence for linkage on chromosome 20 in 716 Finnish affected sib pairs. *Proc Natl Acad Sci U S A* 96:2198-203
- Ghosh S, Watanabe RM, Valle TT, Hauser ER, Magnuson VL, Langefeld CD, Ally DS, et al. (2000) The Finland-United States investigation of non-insulin-dependent diabetes mellitus genetics (FUSION) study. I. An autosomal genome scan for genes that predispose to type 2 diabetes. *Am J Hum Genet* 67:1174-85
- Glazier AM, Nadeau JH, Aitman TJ (2002) Finding genes that underlie complex traits. *Science* 298:2345-9
- Goldstein JL, Schrott HG, Hazzard WR, Bierman EL, Motulsky AG (1973) Hyperlipidemia in coronary heart disease. II. Genetic analysis of lipid levels in 176 families and delineation of a new inherited disorder, combined hyperlipidemia. *J Clin Invest* 52:1544-68
- Gordon DJ, Probstfield JL, Garrison RJ, Neaton JD, Castelli WP, Knoke JD, Jacobs DR, Jr., Bangdiwala S, Tyroler HA (1989) High-density lipoprotein cholesterol and cardiovascular disease. Four prospective American studies. *Circulation* 79:8-15
- Gordon T, Castelli WP, Hjortland MC, Kannel WB, Dawber TR (1977) High density lipoprotein as a protective factor against coronary heart disease. The Framingham Study. *Am J Med* 62:707-14
- Gotto AM, Jr., Pownall HJ, Havel RJ (1986) Introduction to the plasma lipoproteins. *Methods Enzymol* 128:3-41
- Groenendijk M, Cantor RM, Blom NH, Rotter JI, de Bruin TW, Dallinga-Thie GM (1999) Association of plasma lipids and apolipoproteins with the insulin response element in the apoC-III promoter region in familial combined hyperlipidemia. *J Lipid Res* 40:1036-44
- Groenendijk M, Cantor RM, de Bruin TW, Dallinga-Thie GM (2001a) The apoAI-CIII-AIV gene cluster. *Atherosclerosis* 157:1-11
- Groenendijk M, Cantor RM, De Bruin TW, Dallinga-Thie GM (2001b) New genetic variants in the apoA-I and apoC-III genes and familial combined hyperlipidemia. *J Lipid Res* 42:188-94
- Groenendijk M, Cantor RM, Funke H, Dallinga-Thie GM (2001c) Two newly identified SNPs in the APO AI-CIII intergenic region are strongly associated with familial combined hyperlipidaemia. *Eur J Clin Invest* 31:852-9
- Groenendijk M, De Bruin TW, Dallinga-Thie GM (2001d) Two polymorphisms in the apo A-IV gene and familial combined hyperlipidemia. *Atherosclerosis* 158:369-76
- Grundy SM (1999) Hypertriglyceridemia, insulin resistance, and the metabolic syndrome. *Am J Cardiol* 83:25F-29F
- Guerra R, Wang J, Grundy SM, Cohen JC (1997) A hepatic lipase (LIPC) allele associated with high plasma concentrations of high density lipoprotein cholesterol. *Proc Natl Acad Sci U S A* 94:4532-7
- Haffner SM, Miettinen H, Stern MP (1997) The homeostasis model in the San Antonio Heart Study. *Diabetes Care* 20:1087-92

- Hager J, Dina C, Francke S, Dubois S, Houari M, Vatin V, Vaillant E, Lorentz N, Basdevant A, Clement K, Guy-Grand B, Froguel P (1998) A genome-wide scan for human obesity genes reveals a major susceptibility locus on chromosome 10. *Nat Genet* 20:304-8
- Hamsten A, Iselius L, Dahlen G, de Faire U (1986) Genetic and cultural inheritance of serum lipids, low and high density lipoprotein cholesterol and serum apolipoproteins A-I, A-II and B. *Atherosclerosis* 60:199-208
- Han Z, Heath SC, Shmulewitz D, Li W, Auerbach SB, Blundell ML, Lehner T, Ott J, Stoffel M, Friedman JM, Breslow JL (2002) Candidate genes involved in cardiovascular risk factors by a family-based association study on the island of Kosrae, Federated States of Micronesia. *Am J Med Genet* 110:234-42
- Hanson RL, Ehm MG, Pettitt DJ, Prochazka M, Thompson DB, Timberlake D, Foroud T, Kobes S, Baier L, Burns DK, Almasy L, Blangero J, Garvey WT, Bennett PH, Knowler WC (1998) An autosomal genomic scan for loci linked to type II diabetes mellitus and body-mass index in Pima Indians. *Am J Hum Genet* 63:1130-8
- Haseman JK, Elston RC (1972) The investigation of linkage between a quantitative trait and a marker locus. *Behav Genet* 2:3-19
- Hayden MR, Kirk H, Clark C, Frohlich J, Rabkin S, McLeod R, Hewitt J (1987) DNA polymorphisms in and around the Apo-A1-CIII genes and genetic hyperlipidemias. *Am J Hum Genet* 40:421-30
- Hedrick CC, Castellani LW, Warden CH, Puppione DL, Lusis AJ (1993) Influence of mouse apolipoprotein A-II on plasma lipoproteins in transgenic mice. *J Biol Chem* 268:20676-82
- Hinney A, Ziegler A, Oeffner F, Wedewardt C, Vogel M, Wulfstange H, Geller F, Stubing K, Siegfried W, Goldschmidt HP, Remschmidt H, Hebebrand J (2000) Independent confirmation of a major locus for obesity on chromosome 10. *J Clin Endocrinol Metab* 85:2962-5
- Hoffer MJ, Bredie SJ, Boomsma DI, Reymer PW, Kastelein JJ, de Knijff P, Demacker PN, Stalenhoef AF, Havekes LM, Frants RR (1996) The lipoprotein lipase (Asn291-->Ser) mutation is associated with elevated lipid levels in families with familial combined hyperlipidaemia. *Atherosclerosis* 119:159-67
- Hoffer MJ, Bredie SJ, Snieder H, Reymer PW, Demacker PN, Havekes LM, Boomsma DI, Stalenhoef AF, Frants RR, Kastelein JJ (1998) Gender-related association between the -93T-->G/D9N haplotype of the lipoprotein lipase gene and elevated lipid levels in familial combined hyperlipidemia. *Atherosclerosis* 138:91-9
- Hokanson JE, Austin MA, Zambon A, Brunzell JD (1993) Plasma triglyceride and LDL heterogeneity in familial combined hyperlipidemia. *Arterioscler Thromb* 13:427-34
- Howard BV (1987) Lipoprotein metabolism in diabetes mellitus. *J Lipid Res* 28:613-28
- Hsueh WC, St Jean PL, Mitchell BD, Pollin TI, Knowler WC, Ehm MG, Bell CJ, Sakul H, Wagner MJ, Burns DK, Shuldiner AR (2003) Genome-wide and fine-mapping linkage studies of type 2 diabetes and glucose traits in the Old Order Amish: evidence for a new diabetes locus on chromosome 14q11 and confirmation of a locus on chromosome 1q21-q24. *Diabetes* 52:550-7
- Hugot JP, Chamaillard M, Zouali H, Lesage S, Cezard JP, Belaiche J, Almer S, Tysk C, O'Morain CA, Gassull M, Binder V, Finkel Y, Cortot A, Modigliani R, Laurent-Puig P, Gower-Rousseau C, Macry J, Colombel JF, Sahbatou M, Thomas G (2001) Association of NOD2 leucine-rich repeat variants with susceptibility to Crohn's disease. *Nature* 411:599-603
- Hunt SC, Wu LL, Hopkins PN, Stults BM, Kuida H, Ramirez ME, Lalouel JM, Williams RR (1989) Apolipoprotein, low density lipoprotein subfraction, and insulin associations with familial combined hyperlipidemia. Study of Utah patients with familial dyslipidemic hypertension. *Arteriosclerosis* 9:335-44
- Inazu A, Nishimura Y, Terada Y, Mabuchi H (2001) Effects of hepatic lipase gene promoter nucleotide variations on serum HDL cholesterol concentration in the general Japanese population. *J Hum Genet* 46:172-7
- Ilyndjian PB (1998) Identification of upstream stimulatory factor as transcriptional activator of the liver promoter of the glucokinase gene. *Biochem J* 333 (Pt 3):705-12
- Jansen H, Verhoeven AJ, Sijbrands EJ (2002) Hepatic lipase: a pro- or anti-atherogenic protein? *J Lipid Res* 43:1352-62
- Jaworski MA, Severini A, Mansour G, Hennig K, Slater JD, Jeske R, Schlaut J, Yoon JW, Maclaren NK, Nepom GT (1989) Inherited diseases in North American Mennonites: focus on Old Colony (Chortitza) Mennonites. *Am J Med Genet* 32:158-68
- Jeenah M, Kessling A, Miller N, Humphries S (1990) G to A substitution in the promoter region of the apolipoprotein AI gene is associated with elevated serum apolipoprotein AI and high density lipoprotein cholesterol concentrations. *Mol Biol Med* 7:233-41
- Ji L, Malecki M, Warram JH, Yang Y, Rich SS, Krolewski AS (1997) New susceptibility locus for NIDDM is localized to human chromosome 20q. *Diabetes* 46:876-81
- Jiang XC, Qin S, Qiao C, Kawano K, Lin M, Skold A, Xiao X, Tall AR (2001) Apolipoprotein B secretion and atherosclerosis are decreased in mice with phospholipid-transfer protein deficiency. *Nat Med* 7:847-52
- Juo SH, Bredie SJ, Kiemeny LA, Demacker PN, Stalenhoef AF (1998) A common genetic mechanism determines plasma apolipoprotein B levels and dense LDL subfraction distribution in familial combined hyperlipidemia. *Am J Hum Genet* 63:586-94
- Kakko S, Kelloniemi J, von Rohr P, Hoeschele I, Tamminen M, Brousseau ME, Kesaniemi YA, Savolainen MJ (2003) ATP-binding cassette transporter A1 locus is not a major determinant of HDL-C levels in a population at high risk for coronary heart disease. *Atherosclerosis* 166:285-90
- Kannel WB, McGee DL (1979) Diabetes and cardiovascular disease. The Framingham study. *Jama* 241:2035-8

- King H, Aubert RE, Herman WH (1998) Global burden of diabetes, 1995-2025: prevalence, numerical estimates, and projections. *Diabetes Care* 21:1414-31
- Kittles RA, Perola M, Peltonen L, Bergen AW, Aragon RA, Virkkunen M, Linnoila M, Goldman D, Long JC (1998) Dual origins of Finns revealed by Y chromosome haplotype variation. *Am J Hum Genet* 62:1171-9
- Knoblauch H, Busjahn A, Munter S, Nagy Z, Faulhaber HD, Schuster H, Luft FC (1997) Heritability analysis of lipids and three gene loci in twins link the macrophage scavenger receptor to HDL cholesterol concentrations. *Arterioscler Thromb Vasc Biol* 17:2054-60
- Korstanje R, Paigen B (2002) From QTL to gene: the harvest begins. *Nat Genet* 31:235-6
- Kort EN, Ballinger DG, Ding W, Hunt SC, Bowen BR, Abkevich V, Bulka K, Campbell B, Capener C, Gutin A, Harshman K, McDermott M, Thorne T, Wang H, Wardell B, Wong J, Hopkins PN, Skolnick M, Samuels M (2000) Evidence of linkage of familial hypoalphalipoproteinemia to a novel locus on chromosome 11q23. *Am J Hum Genet* 66:1845-56
- Kovanen PT, Pentikainen MO (2003) Circulating lipoproteins as proinflammatory and anti-inflammatory particles in atherogenesis. *Curr Opin Lipidol* 14:411-9
- Krauss RM (1994) Heterogeneity of plasma low-density lipoproteins and atherosclerosis risk. *Curr Opin Lipidol* 5:339-49
- Kruglyak L (1999) Prospects for whole-genome linkage disequilibrium mapping of common disease genes. *Nat Genet* 22:139-44
- Kruglyak L, Nickerson DA (2001) Variation is the spice of life. *Nat Genet* 27:234-6
- Kuivenhoven JA, Pritchard H, Hill J, Frohlich J, Assmann G, Kastelein J (1997) The molecular pathology of lecithin:cholesterol acyltransferase (LCAT) deficiency syndromes. *J Lipid Res* 38:191-205
- Kuokkanen S, Gschwend M, Rioux JD, Daly MJ, Terwilliger JD, Tienari PJ, Wikstrom J, Palo J, Stein LD, Hudson TJ, Lander ES, Peltonen L (1997) Genomewide scan of multiple sclerosis in Finnish multiplex families. *Am J Hum Genet* 61:1379-87
- Kuokkanen S, Sundvall M, Terwilliger JD, Tienari PJ, Wikstrom J, Holmdahl R, Pettersson U, Peltonen L (1996) A putative vulnerability locus to multiple sclerosis maps to 5p14-p12 in a region syntenic to the murine locus Eae2. *Nat Genet* 13:477-80
- Kwiterovich PO, Jr. (2002) Clinical relevance of the biochemical, metabolic, and genetic factors that influence low-density lipoprotein heterogeneity. *Am J Cardiol* 90:30i-47i
- Kwiterovich POJ (1993) Genetics and molecular biology of familial combined hyperlipidemia. *Curr Opin Lipidol* 4:133-143
- Laaksonen DE, Lakka HM, Niskanen LK, Kaplan GA, Salonen JT, Lakka TA (2002) Metabolic syndrome and development of diabetes mellitus: application and validation of recently suggested definitions of the metabolic syndrome in a prospective cohort study. *Am J Epidemiol* 156:1070-7
- Laan M, Paabo S (1997) Demographic history and linkage disequilibrium in human populations. *Nat Genet* 17:435-8
- Lakka HM, Laaksonen DE, Lakka TA, Niskanen LK, Kumpusalo E, Tuomilehto J, Salonen JT (2002) The metabolic syndrome and total and cardiovascular disease mortality in middle-aged men. *Jama* 288:2709-16
- Lander E, Kruglyak L (1995) Genetic dissection of complex traits: guidelines for interpreting and reporting linkage results. *Nat Genet* 11:241-7
- Lander ES, Botstein D (1986) Mapping complex genetic traits in humans: new methods using a complete RFLP linkage map. *Cold Spring Harb Symp Quant Biol* 51 Pt 1:49-62
- Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, Devon K, et al. (2001) Initial sequencing and analysis of the human genome. *Nature* 409:860-921
- Lander ES, Schork NJ (1994) Genetic dissection of complex traits. *Science* 265:2037-48
- Langefeld CD, Wagenknecht LE, Rotter JI, Williams AH, Hokanson JE, Saad MF, Bowden DW, Haffner S, Norris JM, Rich SS, Mitchell BD (2004) Linkage of the metabolic syndrome to 1q23-q31 in Hispanic families: the Insulin Resistance Atherosclerosis Study Family Study. *Diabetes* 53:1170-4
- Lathrop GM, Lalouel JM (1984) Easy calculations of lod scores and genetic risks on small computers. *Am J Hum Genet* 36:460-5
- Lathrop GM, Lalouel JM, Julier C, Ott J (1984) Strategies for multilocus linkage analysis in humans. *Proc Natl Acad Sci U S A* 81:3443-6
- Lathrop GM, Lalouel JM, White RL (1986) Construction of human linkage maps: likelihood calculations for multilocus linkage analysis. *Genet Epidemiol* 3:39-52
- Lee JH, Reed DR, Li WD, Xu W, Joo EJ, Kilker RL, Nanthakumar E, North M, Sakul H, Bell C, Price RA (1999) Genome scan for human obesity and linkage to markers in 20q13. *Am J Hum Genet* 64:196-209
- Lefrancois-Martinez AM, Martinez A, Antoine B, Raymondjean M, Kahn A (1995) Upstream stimulatory factor proteins are major components of the glucose response complex of the L-type pyruvate kinase gene promoter. *J Biol Chem* 270:2640-3
- Leppavuori J, Kujala U, Kinnunen J, Kaprio J, Nissila M, Heliovaara M, Klinger N, Partanen J, Terwilliger JD, Peltonen L (1999) Genome scan for predisposing loci for distal interphalangeal joint osteoarthritis: evidence for a locus on 2q. *Am J Hum Genet* 65:1060-7
- Lewis GF (2002) Lipid metabolism. *Curr Opin Lipidol* 13:97-9

- Li GP, Wang JY, Yan SK, Chen BS, Xue H, Wu G (2004) Genetic effect of two polymorphisms in the apolipoprotein A5 gene and apolipoprotein C3 gene on serum lipids and lipoproteins levels in a Chinese population. *Clin Genet* 65:470-6
- Lohmueller KE, Pearce CL, Pike M, Lander ES, Hirschhorn JN (2003) Meta-analysis of genetic association studies supports a contribution of common variants to susceptibility to common disease. *Nat Genet* 33:177-82
- Mackay TF (2001) The genetic architecture of quantitative traits. *Annu Rev Genet* 35:303-39
- Mahaney MC, Almasy L, Rainwater DL, VandeBerg JL, Cole SA, Hixson JE, Blangero J, MacCluer JW (2003) A quantitative trait locus on chromosome 16q influences variation in plasma HDL-C levels in Mexican Americans. *Arterioscler Thromb Vasc Biol* 23:339-45
- Mamo JC, Proctor SD, Smith D (1998) Retention of chylomicron remnants by arterial tissue; importance of an efficient clearance mechanism from plasma. *Atherosclerosis* 141 Suppl 1:S63-9
- Mar R, Pajukanta P, Allayee H, Groenendijk M, Dallinga-Thie G, Krauss RM, Sinsheimer JS, Cantor RM, de Bruin TW, Lusis AJ (2004) Association of the APOLIPOPROTEIN A1/C3/A4/A5 gene cluster with triglyceride levels and LDL particle size in familial combined hyperlipidemia. *Circ Res* 94:993-9
- Marchesini G, Forlani G, Cerrelli F, Manini R, Natale S, Baraldi L, Ermini G, Savorani G, Zocchi D, Melchionda N (2004) WHO and ATPIII proposals for the definition of the metabolic syndrome in patients with Type 2 diabetes. *Diabet Med* 21:383-7
- Marcil M, Boucher B, Gagne E, Davignon J, Hayden M, Genest J, Jr. (1996) Lack of association of the apolipoprotein A-I-C-III-A-IV gene XmnI and SstI polymorphisms and of the lipoprotein lipase gene mutations in familial combined hyperlipoproteinemia in French Canadian subjects. *J Lipid Res* 37:309-19
- Martin CC, Svitek CA, Oeser JK, Henderson E, Stein R, O'Brien RM (2003a) Upstream stimulatory factor (USF) and neurogenic differentiation/beta-cell E box transactivator 2 (NeuroD/BETA2) contribute to islet-specific glucose-6-phosphatase catalytic-subunit-related protein (IGRP) gene expression. *Biochem J* 371:675-86
- Martin S, Nicaud V, Humphries SE, Talmud PJ (2003b) Contribution of APOA5 gene variants to plasma triglyceride determination and to the response to both fat and glucose tolerance challenges. *Biochim Biophys Acta* 1637:217-25
- Masana L, Ribalta J, Salazar J, Fernandez-Ballart J, Joven J, Cabezas MC (2003) The apolipoprotein AV gene and diurnal triglyceridaemia in normolipidaemic subjects. *Clin Chem Lab Med* 41:517-21
- Mathews CA, Reus VI, Bejarano J, Escamilla MA, Fournier E, Herrera LD, Lowe TL, McInnes LA, Molina J, Ophoff RA, Raventos H, Sandkuijl LA, Service SK, Spesny M, Leon PE, Freimer NB (2004) Genetic studies of neuropsychiatric disorders in Costa Rica: a model for the use of isolated populations. *Psychiatr Genet* 14:13-23
- Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC (1985) Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 28:412-9
- Mehrabian M, Qiao JH, Hyman R, Ruddle D, Laughton C, Lusis AJ (1993) Influence of the apoA-II gene locus on HDL levels and fatty streak development in mice. *Arterioscler Thromb* 13:1-10
- Miettinen HE, Gylling H, Miettinen TA, Viikari J, Paulin L, Kontula K (1997) Apolipoprotein A-I^{Fin}. Dominantly inherited hypoalphalipoproteinemia due to a single base substitution in the apolipoprotein A-I gene. *Arterioscler Thromb Vasc Biol* 17:83-90
- Miettinen HE, Gylling H, Tenhunen J, Virtamo J, Jauhainen M, Huttunen JK, Kantola I, Miettinen TA, Kontula K (1998) Molecular genetic study of Finns with hypoalphalipoproteinemia and hyperalphalipoproteinemia: a novel Gly230 Arg mutation (LCAT^{Fin}) of lecithin:cholesterol acyltransferase (LCAT) accounts for 5% of cases with very low serum HDL cholesterol levels. *Arterioscler Thromb Vasc Biol* 18:591-8
- Miller M, Rhyne J, Hamlette S, Birnbaum J, Rodriguez A (2003) Genetics of HDL regulation in humans. *Curr Opin Lipidol* 14:273-9
- Mohlke KL, Lange EM, Valle TT, Ghosh S, Magnuson VL, Silander K, Watanabe RM, Chines PS, Bergman RN, Tuomilehto J, Collins FS, Boehnke M (2001) Linkage disequilibrium between microsatellite markers extends beyond 1 cM on chromosome 20 in Finns. *Genome Res* 11:1221-6
- Morton NE (1955) Sequential tests for the detection of linkage. *Am J Hum Genet* 7:277-318
- Nabholz CE, von Overbeck J (2004) Gene-environment interactions and the complexity of human genetic diseases. *J Insur Med* 36:47-53
- Nadeau JH (2003) Modifier genes and protective alleles in humans and mice. *Curr Opin Genet Dev* 13:290-5
- Naoumova RP, Bonney SA, Eichenbaum-Voline S, Patel HN, Jones B, Jones EL, Amey J, Colilla S, Neuwirth CK, Allotey R, Seed M, Betteridge DJ, Galton DJ, Cox NJ, Bell GI, Scott J, Shoulders CC (2003) Confirmed locus on chromosome 11p and candidate loci on 6q and 8p for the triglyceride and cholesterol traits of combined hyperlipidemia. *Arterioscler Thromb Vasc Biol* 23:2070-7
- Nardone J, Lee DU, Ansel KM, Rao A (2004) Bioinformatics for the 'bench biologist': how to find regulatory regions in genomic DNA. *Nat Immunol* 5:768-74
- Navab M, Berliner JA, Subbanagounder G, Hama S, Lusis AJ, Castellani LW, Reddy S, Shih D, Shi W, Watson AD, Van Lenten BJ, Vora D, Fogelman AM (2001) HDL and the inflammatory response induced by LDL-derived oxidized phospholipids. *Arterioscler Thromb Vasc Biol* 21:481-8

- NCEP (2001) Executive Summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *Jama* 285:2486-97
- Neel JV (1962) Diabetes mellitus: a "thrifty" genotype rendered detrimental by "progress"? *Am J Hum Genet* 14:353-62
- Nevanlinna HR (1972) The Finnish population structure. A genetic and genealogical study. *Hereditas* 71:195-236
- Nikkila EA, Aro A (1973) Family study of serum lipids and lipoproteins in coronary heart-disease. *Lancet* 1:954-9
- Nobrega M, Pennacchio LA (2003) Comparative Genomic Analysis as a Tool for Biological Discovery. *J Physiol*
- Norio R (2003a) Finnish Disease Heritage I: characteristics, causes, background. *Hum Genet* 112:441-56
- Norio R (2003b) Finnish Disease Heritage II: population prehistory and genetic roots of Finns. *Hum Genet* 112:457-69
- Norio R, Nevanlinna HR, Perheentupa J (1973) Hereditary diseases in Finland; rare flora in rare soul. *Ann Clin Res* 5:109-41
- Norum RA, Lakier JB, Goldstein S, Angel A, Goldberg RB, Block WD, Noffze DK, Dolphin PJ, Edelglass J, Bogorad DD, Alaupovic P (1982) Familial deficiency of apolipoproteins A-I and C-III and precocious coronary-artery disease. *N Engl J Med* 306:1513-9
- Nowak M, Helleboid-Chapman A, Jakel H, Rommens C, Martin G, Duran-Sandoval D, Staels B, Rubin EM, Pennacchio LA, Taskinen MR, Fruchart-Najib J, Fruchart JC (2004) Phosphorylation-dependent down-regulation of apolipoprotein A5 by insulin. Submitted
- Ober C, Abney M, McPeck MS (2001) The genetic dissection of complex traits in a founder population. *Am J Hum Genet* 69:1068-79
- Ogura Y, Bonen DK, Inohara N, Nicolae DL, Chen FF, Ramos R, Britton H, Moran T, Karaliuskas R, Duerr RH, Achkar JP, Brant SR, Bayless TM, Kirschner BS, Hanauer SB, Nunez G, Cho JH (2001) A frameshift mutation in NOD2 associated with susceptibility to Crohn's disease. *Nature* 411:603-6
- Ohman M, Oksanen L, Kaprio J, Koskenvuo M, Mustajoki P, Rissanen A, Salmi J, Kontula K, Peltonen L (2000) Genome-wide scan of obesity in Finnish sibpairs reveals linkage to chromosome Xq24. *J Clin Endocrinol Metab* 85:3183-90
- Ordovas JM, Cassidy DK, Civeira F, Bisgaier CL, Schaefer EJ (1989) Familial apolipoprotein A-I, C-III, and A-IV deficiency and premature atherosclerosis due to deletion of a gene complex on chromosome 11. *J Biol Chem* 264:16339-42
- Ordovas JM, Schaefer EJ, Salem D, Ward RH, Glueck CJ, Vergani C, Wilson PW, Karathanasis SK (1986) Apolipoprotein A-I gene polymorphism associated with premature coronary artery disease and familial hypoalphalipoproteinemia. *N Engl J Med* 314:671-7
- Ott J (1991) Analysis of human genetic linkage. Johns Hopkins University Press, Baltimore
- Packard CJ, Demant T, Stewart JP, Bedford D, Caslake MJ, Schwertfeger G, Bedynek A, Shepherd J, Seidel D (2000) Apolipoprotein B metabolism and the distribution of VLDL and LDL subfractions. *J Lipid Res* 41:305-18
- Pajukanta P, Allayee H, Krass KL, Kuraishy A, Soro A, Lilja HE, Mar R, Taskinen MR, Nuotio I, Laakso M, Rotter JJ, de Bruin TW, Cantor RM, Lusk AJ, Peltonen L (2003) Combined analysis of genome scans of Dutch and Finnish families reveals a susceptibility locus for high-density lipoprotein cholesterol on chromosome 16q. *Am J Hum Genet* 72:903-17
- Pajukanta P, Bodnar JS, Sallinen R, Chu M, Airaksinen T, Xiao Q, Castellani LW, Sheth SS, Wessman M, Palotie A, Sinsheimer JS, Demant P, Lusk AJ, Peltonen L (2001) Fine mapping of Hyplip1 and the human homolog, a potential locus for FCHL. *Mamm Genome* 12:238-45
- Pajukanta P, Cargill M, Viitanen L, Nuotio I, Kareinen A, Perola M, Terwilliger JD, Kempas E, Daly M, Lilja H, Rioux JD, Bretin T, Viikari JS, Ronnema T, Laakso M, Lander ES, Peltonen L (2000) Two loci on chromosomes 2 and X for premature coronary heart disease identified in early- and late-settlement populations of Finland. *Am J Hum Genet* 67:1481-93
- Pajukanta P, Lilja HE, Sinsheimer JS, Cantor RM, Lusk AJ, Gentile M, Duan XJ, Soro-Paavonen A, Naukkarinen J, Saarela J, Laakso M, Ehnholm C, Taskinen MR, Peltonen L (2004) Familial combined hyperlipidemia is associated with upstream transcription factor 1 (USF1). *Nat Genet* 36:371-6
- Pajukanta P, Nuotio I, Terwilliger JD, Porkka KV, Ylitalo K, Pihlajamaki J, Suomalainen AJ, Syvanen AC, Lehtimäki T, Viikari JS, Laakso M, Taskinen MR, Ehnholm C, Peltonen L (1998) Linkage of familial combined hyperlipidaemia to chromosome 1q21-q23. *Nat Genet* 18:369-73
- Pajukanta P, Porkka KV, Antikainen M, Taskinen MR, Perola M, Murtomäki-Repo S, Ehnholm S, Nuotio I, Suurinkero L, Lahdenkari AT, Syvanen AC, Viikari JS, Ehnholm C, Peltonen L (1997) No evidence of linkage between familial combined hyperlipidemia and genes encoding lipolytic enzymes in Finnish families. *Arterioscler Thromb Vasc Biol* 17:841-50
- Pajukanta P, Terwilliger JD, Perola M, Hiekkalinna T, Nuotio I, Ellonen P, Parkkonen M, Hartiala J, Ylitalo K, Pihlajamaki J, Porkka K, Laakso M, Viikari J, Ehnholm C, Taskinen MR, Peltonen L (1999) Genomewide scan for familial combined hyperlipidemia genes in Finnish families, suggesting multiple susceptibility loci influencing triglyceride, cholesterol, and apolipoprotein B levels. *Am J Hum Genet* 64:1453-63
- Pan L, Black TA, Shi Q, Jones CA, Petrovic N, Loudon J, Kane C, Sigmund CD, Gross KW (2001) Critical roles of a cyclic AMP responsive element and an E-box in regulation of mouse renin gene expression. *J Biol Chem* 276:45530-8

- Park YW, Zhu S, Palaniappan L, Heshka S, Carnethon MR, Heymsfield SB (2003) The metabolic syndrome: prevalence and associated risk factor findings in the US population from the Third National Health and Nutrition Examination Survey, 1988-1994. *Arch Intern Med* 163:427-36
- Pastier D, Lacorte JM, Chambaz J, Cardot P, Ribeiro A (2002) Two initiator-like elements are required for the combined activation of the human apolipoprotein C-III promoter by upstream stimulatory factor and hepatic nuclear factor-4. *J Biol Chem* 277:15199-206
- Pastinen T, Perola M, Ignatius J, Sabatti C, Tainola P, Levander M, Syvanen AC, Peltonen L (2001) Dissecting a population genome for targeted screening of disease mutations. *Hum Mol Genet* 10:2961-72
- Patil N, Berno AJ, Hinds DA, Barrett WA, Doshi JM, Hacker CR, Kautzer CR, Lee DH, Marjoribanks C, McDonough DP, Nguyen BT, Norris MC, Sheehan JB, Shen N, Stern D, Stokowski RP, Thomas DJ, Trulson MO, Vyas KR, Frazer KA, Fodor SP, Cox DR (2001) Blocks of limited haplotype diversity revealed by high-resolution scanning of human chromosome 21. *Science* 294:1719-23
- Paunio T, Ekelund J, Varilo T, Parker A, Hovatta I, Turunen JA, Rinard K, Foti A, Terwilliger JD, Juvonen H, Suvisaari J, Arajärvi R, Suokas J, Partonen T, Lonnqvist J, Meyer J, Peltonen L (2001) Genome-wide scan in a nationwide study sample of schizophrenia families in Finland reveals susceptibility loci on chromosomes 2q and 5q. *Hum Mol Genet* 10:3037-48
- Pei W, Baron H, Muller-Miyshok B, Knoblauch H, Al-Yahyaee SA, Hui R, Wu X, Liu L, Busjahn A, Luft FC, Schuster H (2000) Support for linkage of familial combined hyperlipidemia to chromosome 1q21-q23 in Chinese and German families. *Clin Genet* 57:29-34
- Peltonen L, Jalanko A, Varilo T (1999) Molecular genetics of the Finnish disease heritage. *Hum Mol Genet* 8:1913-23
- Peltonen L, McKusick VA (2001) Genomics and medicine. Dissecting human disease in the postgenomic era. *Science* 291:1224-9
- Peltonen L, Palotie A, Lange K (2000) Use of population isolates for mapping complex traits. *Nat Rev Genet* 1:182-90
- Peltonen L, Pekkarinen P, Aaltonen J (1995) Messages from an isolate: lessons from the Finnish gene pool. *Biol Chem Hoppe Seyler* 376:697-704
- Pennacchio LA (2003) Insights from human/mouse genome comparisons. *Mamm Genome* 14:429-36
- Pennacchio LA, Olivier M, Hubacek JA, Cohen JC, Cox DR, Fruchart JC, Krauss RM, Rubin EM (2001) An apolipoprotein influencing triglycerides in humans and mice revealed by comparative sequencing. *Science* 294:169-73
- Pennacchio LA, Rubin EM (2001) Genomic strategies to identify mammalian regulatory sequences. *Nat Rev Genet* 2:100-9
- Pennacchio LA, Rubin EM (2003a) Apolipoprotein A5, a newly identified gene that affects plasma triglyceride levels in humans and mice. *Arterioscler Thromb Vasc Biol* 23:529-34
- Pennacchio LA, Rubin EM (2003b) Comparative genomic tools and databases: providing insights into the human genome. *J Clin Invest* 111:1099-106
- Pennisi E (2003) Human genome. A low number wins the GeneSweep Pool. *Science* 300:1484
- Permutt MA, Wasson JC, Suarez BK, Lin J, Thomas J, Meyer J, Lewitzky S, Rennich JS, Parker A, DuPrat L, Maruti S, Chayen S, Glaser B (2001) A genome scan for type 2 diabetes susceptibility loci in a genetically isolated population. *Diabetes* 50:681-5
- Perola M, Kainulainen K, Pajukanta P, Terwilliger JD, Hiekkalinna T, Ellonen P, Kaprio J, Koskenvuo M, Kontula K, Peltonen L (2000) Genome-wide scan of predisposing loci for increased diastolic blood pressure in Finnish siblings. *J Hypertens* 18:1579-85
- Pihlajamäki J, Karjalainen L, Karhapää P, Vauhkonen I, Laakso M (2000) Impaired free fatty acid suppression during hyperinsulinemia is a characteristic finding in familial combined hyperlipidemia, but insulin resistance is observed only in hypertriglyceridemic patients. *Arterioscler Thromb Vasc Biol* 20:164-70
- Pihlajamäki J, Rissanen J, Heikkinen S, Karjalainen L, Laakso M (1997) Codon 54 polymorphism of the human intestinal fatty acid binding protein 2 gene is associated with dyslipidemias but not with insulin resistance in patients with familial combined hyperlipidemia. *Arterioscler Thromb Vasc Biol* 17:1039-44
- Pimstone SN, Gagne SE, Gagne C, Lupien PJ, Gaudet D, Williams RR, Kotze M, Reymer PW, Defesche JC, Kastelein JJ, et al. (1995) Mutations in the gene for lipoprotein lipase. A cause for low HDL cholesterol levels in individuals heterozygous for familial hypercholesterolemia. *Arterioscler Thromb Vasc Biol* 15:1704-12
- Porkka KV, Viikari JS, Ronnemaa T, Marniemi J, Akerblom HK (1994) Age and gender specific serum lipid and apolipoprotein fractions of Finnish children and young adults. The Cardiovascular Risk in Young Finns Study. *Acta Paediatr* 83:838-48
- Portois L, Tastenoy M, Viollet B, Svoboda M (2002) Functional analysis of the glucose response element of the rat glucagon receptor gene in insulin-producing INS-1 cells. *Biochim Biophys Acta* 1574:175-86
- Prenger VL, Beatty TH, Kwiterovich PO (1992) Genetic determination of high-density lipoprotein-cholesterol and apolipoprotein A-I plasma levels in a family study of cardiac catheterization patients. *Am J Hum Genet* 51:1047-57
- Price RA, Li WD, Bernstein A, Crystal A, Golding EM, Weisberg SJ, Zuckerman WA (2001) A locus affecting obesity in human chromosome region 10p12. *Diabetologia* 44:363-6

- Putt W, Palmen J, Nicaud V, Tregouet DA, Tahri-Daizadeh N, Flavell DM, Humphries SE, Talmud PJ (2004) Variation in USF1 shows haplotype effects, gene:gene and gene:environment associations with glucose and lipid parameters in the European Atherosclerosis Research Study II. *Hum Mol Genet*
- Rader DJ (2003) Regulation of reverse cholesterol transport and clinical implications. *Am J Cardiol* 92:42J-49J
- Rader DJ, Ikewaki K, Duverger N, Feuerstein I, Zech L, Connor W, Brewer HB, Jr. (1993a) Very low high-density lipoproteins without coronary atherosclerosis. *Lancet* 342:1455-8
- Rader DJ, Schaefer JR, Lohse P, Ikewaki K, Thomas F, Harris WA, Zech LA, Dujovne CA, Brewer HB, Jr. (1993b) Increased production of apolipoprotein A-I associated with elevated plasma levels of high-density lipoproteins, apolipoprotein A-I, and lipoprotein A-I in a patient with familial hyperalphalipoproteinemia. *Metabolism* 42:1429-34
- Read ML, Clark AR, Docherty K (1993) The helix-loop-helix transcription factor USF (upstream stimulating factor) binds to a regulatory sequence of the human insulin gene enhancer. *Biochem J* 295 (Pt 1):233-7
- Reed DR, Nanthakumar E, North M, Bell C, Price RA (2001) A genome-wide scan suggests a locus on chromosome 1q21-q23 contributes to normal variation in plasma cholesterol concentration. *J Mol Med* 79:262-9
- Reich DE, Cargill M, Bolk S, Ireland J, Sabeti PC, Richter DJ, Lavery T, Kouyoumjian R, Farhadian SF, Ward R, Lander ES (2001) Linkage disequilibrium in the human genome. *Nature* 411:199-204
- Reich DE, Lander ES (2001) On the allelic spectrum of human disease. *Trends Genet* 17:502-10
- Reymer PW, Gagne E, Groenemeyer BE, Zhang H, Forsyth I, Jansen H, Seidell JC, Kromhout D, Lie KE, Kastelein J, et al. (1995a) A lipoprotein lipase mutation (Asn291Ser) is associated with reduced HDL cholesterol levels in premature atherosclerosis. *Nat Genet* 10:28-34
- Reymer PW, Groenemeyer BE, Gagne E, Miao L, Appelman EE, Seidel JC, Kromhout D, Bijvoet SM, van de Oever K, Bruin T, et al. (1995b) A frequently occurring mutation in the lipoprotein lipase gene (Asn291Ser) contributes to the expression of familial combined hyperlipidemia. *Hum Mol Genet* 4:1543-9
- Ribalta J, Figuera L, Fernandez-Ballart J, Vilella E, Castro Cabezas M, Masana L, Joven J (2002) Newly identified apolipoprotein A-I gene predisposes to high plasma triglycerides in familial combined hyperlipidemia. *Clin Chem* 48:1597-600
- Ribeiro A, Pastier D, Kardassios D, Chambaz J, Cardot P (1999) Cooperative binding of upstream stimulatory factor and hepatic nuclear factor 4 drives the transcription of the human apolipoprotein A-II gene. *J Biol Chem* 274:1216-25
- Rigotti A, Trigatti BL, Penman M, Rayburn H, Herz J, Krieger M (1997) A targeted mutation in the murine gene encoding the high density lipoprotein (HDL) receptor scavenger receptor class B type I reveals its key role in HDL metabolism. *Proc Natl Acad Sci U S A* 94:12610-5
- Risch N, Merikangas K (1996) The future of genetic studies of complex human diseases. *Science* 273:1516-7
- Risch NJ (2000) Searching for genetic determinants in the new millennium. *Nature* 405:847-56
- Rose G, Blackburn H, Gillum R (1982) Cardiovascular survey methods, 2nd ed. World Health Organisation.
- Rose HG, Kranz P, Weinstock M, Juliano J, Haft JI (1973) Inheritance of combined hyperlipoproteinemia: evidence for a new lipoprotein phenotype. *Am J Med* 54:148-60
- Rust S, Rosier M, Funke H, Real J, Amoura Z, Piette JC, Deleuze JF, Brewer HB, Duverger N, Deneffe P, Assmann G (1999) Tangier disease is caused by mutations in the gene encoding ATP-binding cassette transporter 1. *Nat Genet* 22:352-5
- Safeer RS, Cornell MO (2000) The emerging role of HDL cholesterol. Is it time to focus more energy on raising high-density lipoprotein levels? *Postgrad Med* 108:87-90, 93-8
- Salero E, Gimenez C, Zafra F (2003) Identification of a non-canonical E-box motif as a regulatory element in the proximal promoter region of the apolipoprotein E gene. *Biochem J* 370:979-86
- Salisbury BA, Pungliya M, Choi JY, Jiang R, Sun XJ, Stephens JC (2003) SNP and haplotype variation in the human genome. *Mutat Res* 526:53-61
- Schaap FG, Rensen PC, Voshol PJ, Vriens C, Van Der Vliet HN, Chamuleau RA, Havekes LM, Groen AK, Willems Van Dijk K (2004) ApoAV reduces plasma triglycerides by inhibiting very low density lipoprotein-triglyceride (VLDL-TG) production and stimulating lipoprotein lipase-mediated VLDL-TG hydrolysis. *J Biol Chem*
- Schaefer EJ, Genest JJ, Jr., Ordovas JM, Salem DN, Wilson PW (1994) Familial lipoprotein disorders and premature coronary artery disease. *Atherosclerosis* 108 Suppl:S41-54
- Schaefer EJ, Ordovas JM, Law SW, Ghiselli G, Kashyap ML, Srivastava LS, Heaton WH, Albers JJ, Connor WE, Lindgren FT, et al. (1985) Familial apolipoprotein A-I and C-III deficiency, variant II. *J Lipid Res* 26:1089-101
- Scott J, Knott TJ, Priestley LM, Robertson ME, Mann DV, Kostner G, Miller GJ, Miller NE (1985) High-density lipoprotein composition is altered by a common DNA polymorphism adjacent to apoprotein AII gene in man. *Lancet* 1:771-3
- Seda O, Sedova L (2003) New apolipoprotein A-V: comparative genomics meets metabolism. *Physiol Res* 52:141-6
- Shoulders CC (2004) USF1 on trial. *Nat Genet* 36:322-3
- Slatkin M (1994) Linkage disequilibrium in growing and stable populations. *Genetics* 137:331-6
- Smih F, Rouet P, Lucas S, Mairal A, Sengenès C, Lafontan M, Vaulont S, Casado M, Langin D (2002) Transcriptional regulation of adipocyte hormone-sensitive lipase by glucose. *Diabetes* 51:293-300

- Sniderman A, Shapiro S, Marpole D, Skinner B, Teng B, Kwiterovich PO, Jr. (1980) Association of coronary atherosclerosis with hyperapobetalipoproteinemia [increased protein but normal cholesterol levels in human plasma low density (beta) lipoproteins]. *Proc Natl Acad Sci U S A* 77:604-8
- Soro A, Jauhiainen M, Ehnholm C, Taskinen MR (2003) Determinants of low HDL levels in familial combined hyperlipidemia. *J Lipid Res* 44:1536-44
- Soro A, Pajukanta P, Lilja HE, Ylitalo K, Hiekkalinna T, Perola M, Cantor RM, Viikari JS, Taskinen MR, Peltonen L (2002) Genome scans provide evidence for low-HDL-C loci on chromosomes 8q23, 16q24.1-24.2, and 20q13.11 in Finnish families. *Am J Hum Genet* 70:1333-40
- Spielman RS, McGinnis RE, Ewens WJ (1993) Transmission test for linkage disequilibrium: the insulin gene region and insulin-dependent diabetes mellitus (IDDM). *Am J Hum Genet* 52:506-16
- Su Z, Zhang S, Nebert DW, Zhang L, Huang D, Hou Y, Liao L, Xiao C (2002) A novel allele in the promoter of the hepatic lipase is associated with increased concentration of HDL-C and decreased promoter activity. *J Lipid Res* 43:1595-601
- Symula DJ, Frazer KA, Ueda Y, Deneffe P, Stevens ME, Wang ZE, Locksley R, Rubin EM (1999) Functional screening of an asthma QTL in YAC transgenic mice. *Nat Genet* 23:241-4
- Tahvanainen E, Pajukanta P, Porkka K, Nieminen S, Ikavalko L, Nuotio I, Taskinen MR, Peltonen L, Ehnholm C (1998) Haplotypes of the ApoA-I/C-III/A-IV gene cluster and familial combined hyperlipidemia. *Arterioscler Thromb Vasc Biol* 18:1810-7
- Talmud PJ, Hawe E, Martin S, Olivier M, Miller GJ, Rubin EM, Pennacchio LA, Humphries SE (2002) Relative contribution of variation within the APOC3/A4/A5 gene cluster in determining plasma triglycerides. *Hum Mol Genet* 11:3039-46
- Taskinen MR (2003) LDL-cholesterol, HDL-cholesterol or triglycerides--which is the culprit? *Diabetes Res Clin Pract* 61 Suppl 1:S19-26
- Taskinen MR, Kuusi T, Helve E, Nikkila EA, Yki-Jarvinen H (1988) Insulin therapy induces antiatherogenic changes of serum lipoproteins in noninsulin-dependent diabetes. *Arteriosclerosis* 8:168-77
- Terwilliger JD, Goring HH (2000) Gene mapping in the 20th and 21st centuries: statistical methods, data analysis, and experimental design. *Hum Biol* 72:63-132
- Terwilliger JD, Ott J (1992) A haplotype-based 'haplotype relative risk' approach to detecting allelic associations. *Hum Hered* 42:337-46
- Terwilliger JD, Zollner S, Laan M, Paabo S (1998) Mapping genes through the use of linkage disequilibrium generated by genetic drift: 'drift mapping' in small populations with no demographic expansion. *Hum Hered* 48:138-54
- Travers MT, Vallance AJ, Gourlay HT, Gill CA, Klein I, Bottema CB, Barber MC (2001) Promoter I of the ovine acetyl-CoA carboxylase- α gene: an E-box motif at -114 in the proximal promoter binds upstream stimulatory factor (USF)-1 and USF-2 and acts as an insulin-response sequence in differentiating adipocytes. *Biochem J* 359:273-84
- Wadham C, Albanese N, Roberts J, Wang L, Bagley CJ, Gamble JR, Rye KA, Barter PJ, Vadas MA, Xia P (2004) High-density lipoproteins neutralize C-reactive protein proinflammatory activity. *Circulation* 109:2116-22
- Vakkilainen J, Porkka KV, Nuotio I, Pajukanta P, Suurinkeroinen L, Ylitalo K, Viikari JS, Ehnholm C, Taskinen MR (1998) Glucose intolerance in familial combined hyperlipidaemia. EUFAM study group. *Eur J Clin Invest* 28:24-32
- Wall JD, Pritchard JK (2003) Haplotype blocks and linkage disequilibrium in the human genome. *Nat Rev Genet* 4:587-97
- Vallet VS, Casado M, Henrion AA, Bucchini D, Raymondjean M, Kahn A, Vaulont S (1998) Differential roles of upstream stimulatory factors 1 and 2 in the transcriptional response of liver genes to glucose. *J Biol Chem* 273:20175-9
- van der Kallen CJ, Cantor RM, van Greevenbroek MM, Geurts JM, Bouwman FG, Aouizerat BE, Allayee H, Buurman WA, Lusis AJ, Rotter JI, de Bruin TW (2000) Genome scan for adiposity in Dutch dyslipidemic families reveals novel quantitative trait loci for leptin, body mass index and soluble tumor necrosis factor receptor superfamily 1A. *Int J Obes Relat Metab Disord* 24:1381-91
- van der Vleuten GM, Hijmans A, Kluijtmans LA, Blom HJ, Stalenhoef AF, de Graaf J (2004) Thioredoxin interacting protein in Dutch families with familial combined hyperlipidemia. *Am J Med Genet* 130A:73-5
- van der Vliet HN, Schaap FG, Levels JH, Ottenhoff R, Looije N, Wesseling JG, Groen AK, Chamuleau RA (2002) Adenoviral overexpression of apolipoprotein A-V reduces serum levels of triglycerides and cholesterol in mice. *Biochem Biophys Res Commun* 295:1156-9
- Van Eerdekewegh P, Little RD, Dupuis J, Del Mastro RG, Falls K, Simon J, Torrey D, et al. (2002) Association of the ADAM33 gene with asthma and bronchial hyperresponsiveness. *Nature* 418:426-30
- Wang D, Sul HS (1995) Upstream stimulatory factors bind to insulin response sequence of the fatty acid synthase promoter. USF1 is regulated. *J Biol Chem* 270:28716-22
- Wang D, Sul HS (1997) Upstream stimulatory factor binding to the E-box at -65 is required for insulin regulation of the fatty acid synthase promoter. *J Biol Chem* 272:26367-74
- Wang X, Paigen B (2002) Quantitative trait loci and candidate genes regulating HDL cholesterol: a murine chromosome map. *Arterioscler Thromb Vasc Biol* 22:1390-401

- Warden CH, Daluiski A, Bu X, Purcell-Huynh DA, De Meester C, Shieh BH, Puppione DL, Gray RM, Reaven GM, Chen YD, et al. (1993) Evidence for linkage of the apolipoprotein A-II locus to plasma apolipoprotein A-II and free fatty acid levels in mice and humans. *Proc Natl Acad Sci U S A* 90:10886-90
- Vartiainen E, Jousilahti P, Alfthan G, Sundvall J, Pietinen P, Puska P (2000) Cardiovascular risk factor changes in Finland, 1972-1997. *Int J Epidemiol* 29:49-56
- Vartiainen E, Puska P, Jousilahti P, Korhonen HJ, Tuomilehto J, Nissinen A (1994) Twenty-year trends in coronary risk factors in north Karelia and in other areas of Finland. *Int J Epidemiol* 23:495-504
- Watanabe RM, Ghosh S, Langefeld CD, Valle TT, Hauser ER, Magnuson VL, Mohlke KL, et al. (2000) The Finland-United States investigation of non-insulin-dependent diabetes mellitus genetics (FUSION) study. II. An autosomal genome scan for diabetes-related quantitative-trait loci. *Am J Hum Genet* 67:1186-200
- Weissenbach J (2004) Genome sequencing: differences with the relatives. *Nature* 429:353-5
- Venter JC, Adams MD, Myers EW, Li PW, Mural RJ, Sutton GG, Smith HO, et al. (2001) The sequence of the human genome. *Science* 291:1304-51
- WHO (1999) Report of a WHO consultation: Definition, diagnosis and classification of diabetes mellitus and its complications. World Health Organization, Geneva, pp 31-33
- WHO (2003) Report of the joint WHO/FAO expert consultation: Diet, nutrition and the prevention of chronic diseases. World Health Organization, Geneva
- Wilson PW, Abbott RD, Castelli WP (1988) High density lipoprotein cholesterol and mortality. The Framingham Heart Study. *Arteriosclerosis* 8:737-41
- Wiltshire S, Hattersley AT, Hitman GA, Walker M, Levy JC, Sampson M, O'Rahilly S, et al. (2001) A genomewide scan for loci predisposing to type 2 diabetes in a U.K. population (the Diabetes UK Warren 2 Repository): analysis of 573 pedigrees provides independent replication of a susceptibility locus on chromosome 1q. *Am J Hum Genet* 69:553-69
- Viollet B, Lefrancois-Martinez AM, Henrion A, Kahn A, Raymondjean M, Martinez A (1996) Immunochemical characterization and transacting properties of upstream stimulatory factor isoforms. *J Biol Chem* 271:1405-15
- Vionnet N, Hani El H, Dupont S, Gallina S, Francke S, Dotte S, De Matos F, Durand E, Lepretre F, Lecoeur C, Gallina P, Zekiri L, Dina C, Froguel P (2000) Genomewide search for type 2 diabetes-susceptibility genes in French whites: evidence for a novel susceptibility locus for early-onset diabetes on chromosome 3q27-qter and independent replication of a type 2-diabetes locus on chromosome 1q21-q24. *Am J Hum Genet* 67:1470-80
- Wojciechowski AP, Farrall M, Cullen P, Wilson TM, Bayliss JD, Farren B, Griffin BA, Caslake MJ, Packard CJ, Shepherd J, et al. (1991) Familial combined hyperlipidaemia linked to the apolipoprotein AI-CII-AIV gene cluster on chromosome 11q23-q24. *Nature* 349:161-4
- von Eckardstein A, Nofer JR, Assmann G (2001) High density lipoproteins and arteriosclerosis. Role of cholesterol efflux and reverse cholesterol transport. *Arterioscler Thromb Vasc Biol* 21:13-27
- Vu-Dac N, Gervois P, Jakel H, Nowak M, Bauge E, Dehondt H, Staels B, Pennacchio LA, Rubin EM, Fruchart-Najib J, Fruchart JC (2003) Apolipoprotein A5, a crucial determinant of plasma triglyceride levels, is highly responsive to peroxisome proliferator-activated receptor alpha activators. *J Biol Chem* 278:17982-5
- Xiang K, Wang Y, Zheng T, Jia W, Li J, Chen L, Shen K, Wu S, Lin X, Zhang G, Wang C, Wang S, Lu H, Fang Q, Shi Y, Zhang R, Xu J, Weng Q (2004) Genome-wide search for type 2 diabetes/impaired glucose homeostasis susceptibility genes in the Chinese: significant linkage to chromosome 6q21-q23 and chromosome 1q21-q24. *Diabetes* 53:228-34
- Xu CF, Talmud P, Schuster H, Houlston R, Miller G, Humphries S (1994) Association between genetic variation at the APO AI-CIII-AIV gene cluster and familial combined hyperlipidaemia. *Clin Genet* 46:385-97
- Yamakawa-Kobayashi K, Yanagi H, Fukayama H, Hirano C, Shimakura Y, Yamamoto N, Arinami T, Tsuchiya S, Hamaguchi H (1999) Frequent occurrence of hypoalphalipoproteinemia due to mutant apolipoprotein A-I gene in the population: a population-based survey. *Hum Mol Genet* 8:331-6
- Yang XP, Freeman LA, Knapper CL, Amar MJ, Remaley A, Brewer HB, Jr., Santamarina-Fojo S (2002) The E-box motif in the proximal ABCA1 promoter mediates transcriptional repression of the ABCA1 gene. *J Lipid Res* 43:297-306
- Ylisaukko-oja T, Nieminen-von Wendt T, Kempas E, Sarenius S, Varilo T, von Wendt L, Peltonen L, Jarvela I (2004) Genome-wide scan for loci of Asperger syndrome. *Mol Psychiatry* 9:161-8
- Zhong S, Sharp DS, Grove JS, Bruce C, Yano K, Curb JD, Tall AR (1996) Increased coronary heart disease in Japanese-American men with mutation in the cholesteryl ester transfer protein gene despite increased HDL levels. *J Clin Invest* 97:2917-23
- Zouali H, Hani EH, Philippi A, Vionnet N, Beckmann JS, Demeais F, Froguel P (1997) A susceptibility locus for early-onset non-insulin dependent (type 2) diabetes mellitus maps to chromosome 20q, proximal to the phosphoenolpyruvate carboxykinase gene. *Hum Mol Genet* 6:1401-8

PREVIOUSLY PUBLISHED IN THIS SERIES BY THE DEPARTMENT OF MOLECULAR MEDICINE

1. **Irma Järvelä**, Molecular distinction of neuronal ceroid-lipofuscinoses: Assignment of separate gene loci for infantile and juvenile forms. NPHI A2/1991
2. **Päivi Helminen**, Hypervariable regions of human genome applied to paternity testing and detection of malignant cell clones. NPHI A1/1992
3. **Elina Ikonen**, Molecular genetics of aspartylglucosaminuria. NPHI A2/1992
4. **Antti Sajantila**, DNA analysis in forensic medicine: Application of the polymerase chain reaction (PCR) to the identification of individuals. NPHI A3/1992
5. **Katariina Kainulainen**, Molecular genetics of Marfan syndrome. NPHI A4/1992
6. **Raili Kauppinen**, Prognosis of acute porphyrias and molecular genetics of acute intermittent porphyria in Finland. NPHI A8/1992
7. **Miikka Vikkula**, The human type II collagen gene and cartilage diseases. NPHI A1/1993
8. **Anu Suomalainen**, Mutations of mitochondrial DNA in human disease. NPHI A4/1993
9. **Pentti Tienari**, Genetic susceptibility in multiple sclerosis. NPHI A10/1993
10. **Nina Enomaa**, Aspartylglucosaminuria: Molecular pathogenesis and in vitro correction of the enzyme defect. KTL A9/1994
11. **Tiina Paunio**, Molecular pathogenesis of familial amyloidosis, Finnish type. NPHI A5/1995
12. **Jouni Vesa**, The molecular defect in infantile neuronal ceroid lipofuscinosis. KTL A12/1995
13. **Elina Hellsten**, Positional cloning of the infantile neuronal ceroid lipofuscinosis gene. KTL A16/1995
14. **Pekka Nokelainen**, Genetic analyses in myotonic dystrophy and tibial muscular dystrophy in Finland. NPHI A5/1996
15. **Ritva Tikkanen**, Human lysosomal aspartylglucosaminidase: Structure, function and intracellular targeting. KTL A4/1996
16. **Aija Riikonen-Kyttälä**, Intracellular maturation of aspartylglucosaminidase. KTL A7/1996
17. **Leena Karttunen**, Molecular pathogenesis of Marfan syndrome. KTL A9/1996
18. **Johanna Aaltonen**, Molecular genetics of APECED (Autoimmune PolyEndocrinopathy-Candidiasis-Ectodermal Dystrophy). KTL A3/1998
19. **Terhi Rantamäki-Häkkinen**, Fibrillin defects in Marfan syndrome: Impact on DNA diagnosis and molecular pathogenesis. KTL A5/1998
20. **Minna Peltola**, Aspartylglucosaminuria (AGU): Lysosomal targeting of AGA, the cellular consequences of mutations and an attempt at gene therapy in the AGU mouse. NPHI A12/1998
21. **Annukka Uusitalo**, Aspartylglucosaminuria: Disease pathogenesis, developmental expression and regulation of the aspartylglucosaminidase gene. NPHI A15/1998
22. **Satu Kuokkanen**, Search for gene loci predisposing to multiple sclerosis in the Finnish population. NPHI A13/1998
23. **Kaisu Nikali**, Molecular genetics of infantile onset spinocerebellar ataxia. NPHI A14/1998
24. **Lasse Lönnqvist**, Molecular pathology of type-1 fibrillinopathies. KTL A16/1998
25. **Kai Tenhunen**, Mouse aspartylglucosaminidase gene and mouse model for aspartylglucosaminuria. KTL A17/1998
26. **Petra Pekkarinen**, Genetic mapping of the loci for a monogenic and multifactorial neuropsychiatric disorder: PLO-SL and familial bipolar disorder. KTL A19/1998
27. **Iiris Hovatta**, Molecular genetics of familial schizophrenia and PLO-SL. KTL A20/1998
28. **Paulina Paavola**, Molecular genetics of Meckel syndrome. KTL A21/1998
29. **Tuomas Klockars**, Positional cloning of the CLN5 gene. KTL A22/1998
30. **Päivi Pajukanta**, The search for familial combined hyperlipidemia susceptibility genes. KTL A26/1998
31. **Markus Perola**, Molecular genetics of hypertension and related traits. KTL A8/1999
32. **Teppo Varilo**, The age of mutations in the Finnish disease heritage; a genealogical and linkage disequilibrium study. KTL A21/1999
33. **Petra Björnses**, Autoimmune polyendocrinopathy – Candidiasis – Ectodermal Dystrophy (APECED): From locus to defective protein. KTL A24/1999
34. **Minna Savukoski**, Molecular genetics of the late infantile neuronal ceroid lipofuscinosis (LINCL): One gene (CLN5) and two gene loci (CLN2 and CLN6). KTL A25/1999
35. **Jyrki Kaukonen**, Autosomal dominant progressive external ophthalmoplegia (adPEO): A tale of two genomes. KTL A4/2000
36. **Tomi Pastinen**, Scoring human genomic SNPs and mutations: Multiplexed primer extension with manifolds and microarrays as solid-support. KTL A5/2000
37. **Hannele Kangas**, Familial amyloidosis of the Finnish type (FAF) – consequences of amyloidosis-associated mutation for gelsolin processing and function. KTL A9/2000
38. **Miina Öhman**, The search for genes predisposing to obesity. KTL A3/2001
39. **Jesper Ekelund**, Molecular genetics of schizophrenia and comorbid and related traits. KTL A17/2001
40. **Tarja Salonen**, Molecular and cellular biology of infantile neuronal ceroid lipofuscinosis (INCL). KTL

- A16/2001
41. **Sonja Jaari**, Proteins involved in high density lipoprotein metabolism: A special reference to apolipoprotein A-I, hepatic lipase and phospholipid transfer protein. KTL A1/2002
 42. **Mari Auranen**, Molecular genetics of autism spectrum disorders in the Finnish population. KTL A23/2002
 43. **Ilona Visapää**, Molecular genetics of the GRACILE syndrome. KTL A28/2002
 44. **Saara Laitinen**, Family of human oxysterol binding protein homologues: ORP2 is a new regulator of cellular lipid metabolism. KTL A30/2002
 45. **Juha Isosomppi**, Molecular and cell biology of infantile (CLN1) and variant late infantile (CLN5) neuronal ceroid lipofuscinoses. KTL A3/2003
 46. **Maria Halonen**, Monogenic model for autoimmune diseases: Molecular basis of autoimmune polyendocrinopathy - candidiasis - ectodermal dystrophy (APECED). KTL A4/2003
 47. **Juha Paloneva**, Two genes behind PLOSL: Molecular and pathological characteristics of the disease. KTL A8/2003
 48. **Titta S. Blom**, Characterisation of cellular defects in Niemann-Pick type C disease. KTL A11/2003
 49. **Nina Aula**, Molecular pathogenesis of Salla disease. KTL A19/2003
 50. **Henna Haravuori**, Molecular genetics of tibial muscular dystrophy (TMD) and a novel distal myopathy. KTL A24/2003
 51. **Riikka Nissinen**, Immunological features of chronic active rheumatoid arthritis. KTL A22/2003
 52. **Jani Saarela**, Characterization of aspartylglucosaminidase activation and aspartylglucosaminuria mutations. KTL A1/2004
 53. **Ville Holmberg**, CLN5 - from mutation to defective protein and clinical phenotype. KTL A2/2004